

# Risks of later abortions on subsequent births

March 4 2020

---



Credit: CC0 Public Domain

New research indicates that a prior induced abortion poses only a very small risk of negative effects on births from subsequent pregnancies, but the risk is higher if the abortion is performed later in the pregnancy.

The *Acta Obstetricia et Gynecologica Scandinavica* study included 418,690 first-time mothers with singleton births between 1996 and 2013 in Finland. Investigators observed an [increased risk](#) of extremely [preterm birth](#) and very low birthweight among women with late abortion(s) when compared with those with early abortion(s).

When the analysis was restricted to only one abortion, women who had had a late abortion showed a higher risk of only extremely preterm birth compared with women who had experienced an early abortion.

**More information:** Situ KC et al, The duration of gestation at previous induced abortion and its impacts on subsequent births: A nationwide registry-based study, *Acta Obstetricia et Gynecologica Scandinavica* (2020). [DOI: 10.1111/aogs.13788](https://doi.org/10.1111/aogs.13788)

Provided by Wiley

Citation: Risks of later abortions on subsequent births (2020, March 4) retrieved 25 April 2024 from <https://medicalxpress.com/news/2020-03-abortion-subsequent-births.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--