

Study shows low carb diet may prevent, reverse age-related effects within the brain

March 5 2020



Lilianne R. Mujica-Parodi, PhD. Credit: Stony Brook University

A study using neuroimaging led by Stony Brook University professor

and lead author Lilianne R. Mujica-Parodi, Ph.D., and published in PNAS, reveals that neurobiological changes associated with aging can be seen at a much younger age than would be expected, in the late 40s. However, the study also suggests that this process may be prevented or reversed based on dietary changes that involve minimizing the consumption of simple carbohydrates.

To better understand how [diet](#) influences [brain](#) aging, the research team focused on the presymptomatic period during which prevention may be most effective. In the article titled "Diet modulates brain network stability, a biomarker for brain aging, in [young adults](#)," they showed, using large-scale life span neuroimaging datasets, that functional communication between [brain regions](#) destabilizes with age, typically in the late 40's, and that destabilization correlates with poorer cognition and accelerates with insulin resistance. Targeted experiments then showed this biomarker for brain aging to be reliably modulated with consumption of different [fuel](#) sources: glucose decreases, and ketones increase, the stability of brain networks. This effect was replicated across both changes to total diet as well as after drinking a fuel-specific calorie-matched supplement.

"What we found with these experiments involves both bad and good news," said Mujica-Parodi, a Professor in the Department of Biomedical Engineering with joint appointments in the College of Engineering & Applied Sciences and Renaissance School of Medicine at Stony Brook University, and a faculty member in the Laufer Center for Physical and Quantitative Biology. "The bad news is that we see the first signs of brain aging much earlier than was previously thought. However, the good news is that we may be able to prevent or reverse these effects with diet, mitigating the impact of encroaching hypometabolism by exchanging glucose for ketones as fuel for neurons."

What the researchers discovered, using neuroimaging of the brain, is that

quite early on there is breakdown of communication between brain regions ("network stability").

"We think that, as people get older, their brains start to lose the ability to metabolize glucose efficiently, causing neurons to slowly starve, and brain networks to destabilize," said Mujica-Parodi. "Thus, we tested whether giving the brain a more efficient fuel source, in the form of ketones, either by following a low-carb diet or drinking ketone supplements, could provide the brain with greater energy. Even in younger individuals, this added energy further stabilized brain networks."

To conduct their experiments, brain network stability was established as a biomarker for aging by using two large-scale brain neuroimaging (fMRI) datasets totaling nearly 1,000 individuals, ages 18 to 88. Destabilization of brain networks was associated with impaired cognition and was accelerated with Type 2 diabetes, an illness that blocks neurons' ability to effectively metabolize glucose. To identify the mechanism as being specific to energy availability, the researchers then held age constant and scanned an additional 42 adults under the age of 50 years with fMRI. This allowed them to observe directly the impact of glucose and ketones on each individual's brain.

The brain's response to diet was tested in two ways. The first was holistic, comparing brain network stability after participants had spent one week on a standard (unrestricted) vs. low carb (for example: meat or fish with salad, but no sugar, grains, rice, starchy vegetables) diet. In a standard diet, the primary fuel metabolized is glucose, whereas in a low-carb diet, the primary fuel metabolized is ketones. However, there might have been other differences between diets driving the observed effects. Therefore, to isolate glucose vs. ketones as the crucial difference between the diets, an independent set of participants was scanned before and after drinking a small dose of glucose on one day, and ketones on the

other, where the two fuels were individually weight-dosed and calorically matched. The results replicated, showing that the differences between the diets could be attributed to the type of fuel they provide to the brain.

Additional findings from the study included the following: Effects of brain aging emerged at age 47, with most rapid degeneration occurring at age 60. Even in younger adults, under age 50, dietary ketosis (whether achieved after one week of dietary change or 30 minutes after drinking ketones) increased overall brain activity and stabilized functional networks. This is thought to be due to the fact that ketones provide greater energy to cells than glucose, even when the fuels are calorically matched. This benefit has previously been shown for the heart, but the current set of experiments provides the first evidence for equivalent effects in the brain.

"This effect matters because brain aging, and especially dementia, are associated with "hypometabolism," in which neurons gradually lose the ability to effectively use [glucose](#) as fuel. Therefore, if we can increase the amount of energy available to the brain by using a different fuel, the hope is that we can restore the brain to more youthful functioning. In collaboration with Dr. Eva Ratai at Massachusetts General Hospital, we're currently addressing this question, by now extending our studies to older populations," said Mujica-Parodi.

"Additional research with collaborators at Children's National, under the direction of Dr. Nathan Smith, focuses on discovering the precise mechanisms by which fuel impacts signaling between neurons. Finally, in collaboration with Dr. Ken Dill and Dr. Steven Skiena, at Stony Brook, we're working on building a comprehensive computational model that can incorporate our understanding of the biology, from individual neurons to whole brains to cognition, as it develops."

More information: Lilianne R. Mujica-Parodi et al. Diet modulates

brain network stability, a biomarker for brain aging, in young adults, *Proceedings of the National Academy of Sciences* (2020). [DOI: 10.1073/pnas.1913042117](https://doi.org/10.1073/pnas.1913042117)

Provided by Stony Brook University

Citation: Study shows low carb diet may prevent, reverse age-related effects within the brain (2020, March 5) retrieved 13 July 2024 from <https://medicalxpress.com/news/2020-03-carb-diet-reverse-age-related-effects.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.