

Certain combinations of cardiovascular drugs may reduce dementia risk

March 11 2020, by Jenesse Miller



Credit: AI-generated image ([disclaimer](#))

Specific combinations of statins and antihypertensives may also reduce risk for Alzheimer's disease, according to a new USC study of nearly 700,000 Medicare beneficiaries.

The findings suggest that treatments already in use for [blood pressure](#)

and cholesterol control could reduce the number of people with Alzheimer's and related dementias, researchers said.

The study was published in *PLOS One*.

"Our research found [dementia risk](#) may be reduced with a specific combinations of [drug](#) treatments for vascular health," said lead author Julie Zissimopoulos, director of the Aging and Cognition program at the USC Leonard D. Schaeffer Center for Health Policy & Economics.

"This is the first study to investigate the association of different combinations of these frequently used drugs and dementia risk in a large and broadly representative sample of older Americans and to illuminate the differences in effect for women compared to men, and for whites, Hispanics and African Americans."

Dementia affects about 7 million Americans and is projected to grow to 12 million over the next two decades. With no proven treatments, there has been increased global attention on prevention and risk reduction of dementia, including maintenance of a healthy lifestyle and management of risk factors such as high blood pressure and high cholesterol.

Approximately one in four adults over age 65 use both antihypertensives and statins for these conditions.

"We know that managing hyperlipidemia and hypertension is important, and this study tells us there might be certain combinations of drugs that have additional benefits for Alzheimer's and other dementias beyond the management of those targeted conditions," said co-author Douglas Barthold, research assistant professor with the Comparative Health Outcomes, Policy, and Economics (CHOICE) Institute at the Department of Pharmacy at the University of Washington, Seattle.

Specific combinations of drugs may hold the key to

reduced risk

The study examined the medical and pharmacy claims of a random 20% sample of Medicare beneficiaries, aged 67 and higher, enrolled in traditional Medicare from 2007 to 2014. Researchers linked claims from Medicare Parts A (inpatient care), B (outpatient care) and D ([prescription drugs](#)) to enrollment files that included beneficiaries' characteristics such as race and sex.

To be included in the analysis, individuals had to have used both an antihypertensive and a statin for the two previous years and have no prior dementia diagnoses and no prior use of Alzheimer's disease-specific medications.

The study found the use of cholesterol-lowering drugs pravastatin and rosuvastatin, combined with ACE inhibitors or angiotensin II receptor blockers (ARBs) for [high blood pressure](#), was associated with reduced risk for dementia as compared to other combinations of drugs.

The risk was especially lower for people using pravastatin and rosuvastatin in [combination](#) with ARBs, and more so for men than women.

The magnitudes of estimated risk reductions were meaningful; for example, using ARBs combined with pravastatin was associated with 21% lower odds of a dementia diagnosis, as compared to individuals using other combinations of drugs.

"We don't currently have drugs that are proven to treat [dementia](#), but even small delays in onset can dramatically reduce the burden on patients, caregivers and the health system as a whole," Zissimopoulos said.

The study authors say more research is needed but that if their findings are replicated, specific combinations of statins and antihypertensives might be recommended in the interest of reducing the risk of Alzheimer's disease and related dementias.

More information: Douglas Barthold et al. Association of combination statin and antihypertensive therapy with reduced Alzheimer's disease and related dementia risk, *PLOS ONE* (2020). DOI: [10.1371/journal.pone.0229541](https://doi.org/10.1371/journal.pone.0229541)

Provided by University of Southern California

Citation: Certain combinations of cardiovascular drugs may reduce dementia risk (2020, March 11) retrieved 5 May 2024 from <https://medicalxpress.com/news/2020-03-combinations-cardiovascular-drugs-dementia.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|