


Comprehensive COVID-19 hospitalisation and death rate estimates help countries best prepare as global pandemic unfolds

March 31 2020

Estimated proportion of all COVID-19 infections that will require hospitalisation



The proportion of all infections that are hospitalised, estimated from a subset of 3665 cases reported in mainland China up to Feb 8, 2020, and adjusted for demography and under-ascertainment. UK cases classified as 'severe' are assumed to require hospitalisation.

Hospitalisation rates nearly double from 4.3% in 40-49-year-olds to 8.2% in 50-59-year olds
Estimates are constantly being updated and improved over time.

Estimates of the severity of coronavirus disease 2019: a model-based analysis *The Lancet Infectious Diseases*

COVID-19 publication characteristics	Format Research article	Externally peer reviewed? Yes	Date published 30 March 2020
--------------------------------------	----------------------------	----------------------------------	---------------------------------

THE LANCET Infectious Diseases

The best science for better lives

The estimated proportion of hospitalization by age. Credit: The Lancet Infectious Diseases

First comprehensive estimates from mainland China of the proportion of

people with COVID-19 who required hospitalisation, and latest death rate estimates, both show sharp increases with age. Authors warn that without intervention the number of people needing hospital treatment is likely to overwhelm even the most advanced healthcare systems worldwide, though caution that as the pandemic unfolds, it is possible that outcomes could improve and it will be important to revise estimates in this study.

Key findings:

- The death rate from confirmed COVID-19 cases is estimated at 1.38%, while the overall death rate, which includes unconfirmed cases, is estimated at 0.66%; these rates are slightly lower than some estimates for COVID-19 to date, which had not adjusted for undiagnosed cases or for the number of people in each age group of a population
- Death rates vary substantially, ranging from 0.0016% in 0 to 9-year-olds to 7.8% for people aged 80 and above
- Differences in [hospitalisation](#) rates by age were reported, increasing with age—with 11.8% of people in their 60s, 16.6% of people in their 70s, and 18.4% of those in their 80s and above estimated to develop symptoms severe enough for hospitalisation
- These hospitalisation rates compare with 0.04% of 10 to 19-year-olds, 1.0% of people in their 20s, and 3.4% of people aged 30 to 39. Hospitalisation rates nearly double from 4.3% in 40-49-year-olds to 8.2% in 50-59-year olds

Nearly one in five over-80s infected with COVID-19 are likely to require hospitalisation, compared with around 1% of people under 30, according to an analysis of 3,665 cases in mainland China, published in *The Lancet Infectious Diseases* journal.


The new analysis also finds that the estimated proportion of deaths from both diagnosed cases and from milder, unconfirmed cases is strongly influenced by age. The estimates are slightly lower than others that have been made for the virus, but are still much higher than for previous pandemics such as 2009 pandemic influenza H1N1, which was estimated to be fatal in around 0.02% of cases. The new estimates are based on an analysis of 70,117 laboratory-confirmed and clinically-diagnosed cases in mainland China, combined with 689 positive cases among people evacuated from Wuhan on repatriation flights.

"This study provides critical estimates on the proportion of people requiring hospitalisation which, when applied to the UK population, enabled us to get a handle on how many people might need to access NHS services," says Professor Neil Ferguson from Imperial College London, UK. "As the UK epidemic unfolds, more data are becoming available, and at the moment the proportion of people in each age group most likely to require hospitalisation, and most likely to die from infection, are consistent with the estimates in this study."

The authors warn that as 50% to 80% of the global population could be infected with COVID-19 the number of people needing hospital treatment is likely to overwhelm even the most advanced healthcare systems worldwide. However, they caution that it is possible that outcomes could improve, in which case it will be important to revise the estimates in this study.

"Our estimates can be applied to any country to inform decisions around the best containment policies for COVID-19," says Professor Azra Ghani from Imperial College London, UK. "There might be outlying cases that get a lot of media attention, but our analysis very clearly shows that at aged 50 and over, hospitalisation is much more likely than in those under 50, and a greater proportion of cases are likely to be fatal."

Estimated death rates from **confirmed cases*** and from all infections, including undiagnosed with mild symptoms



■ Estimated proportion of confirmed cases* that will die

*Based on 70,117 laboratory-confirmed and clinically-diagnosed cases in mainland China from 1 January to 11 February 2020, with the presented values adjusted for demography and undetected symptomatic cases

■ Estimated proportion of all people infected who die from the disease, including those with only mild symptoms that are not formally diagnosed (estimated by combining confirmed case data in mainland China with information on infection prevalence among those returning home on repatriation flights)

An average of **0.15%** of infected people under 60 are estimated to die

An average of **3.3%** of infected people over 60 are estimated to die

This does not separate out the effect of underlying health conditions, which often correlate with age, and may vary geographically. Outcomes may improve over time, so estimates will need to be updated.

Estimates of the severity of coronavirus disease 2019: a model-based analysis *The Lancet Infectious Diseases*

COVID-19 publication characteristics	Format Research article	Externally peer reviewed? Yes	Date published 30 March 2020
--	-------------------------------	----------------------------------	------------------------------------

THE LANCET Infectious Diseases

The best science for better lives

Estimated death rates by age. Credit: The Lancet Infectious Diseases

Previous estimates of deaths from confirmed cases of COVID-19 have ranged from 2% to 8%, while deaths from overall infections have been estimated at 0.2% to 1.6%. Also, estimates for the proportion of deaths in the oldest age group, the over-80s, have been estimated to be between 8% to 36%. However, these past estimates had not adjusted for the fact that only people with more severe symptoms are likely to be tested, or people in quarantine following repatriation to other countries, so they did not reflect the true number of cases across populations. No previous studies have estimated the proportion of infections that will require hospitalisation.

For the current analysis, a team of international researchers used 3,665 cases from mainland China to estimate the proportion of cases likely to be severe enough to require hospitalisation. To estimate the average time between a person displaying symptoms and dying, they analysed 24 deaths in Hubei Province. The average recovery time was estimated using data from 2,010 international cases, of whom 169 people recovered. Death rates from confirmed cases were estimated using data on 44,672 cases in mainland China. To estimate death rates relevant to the wider population, data from 689 people repatriated from Wuhan to other countries and 3,711 people quarantined on board the cruise liner Diamond Princess were used.

For all the estimates, the researchers assumed that people of all ages are equally likely to become infected, which is consistent with previous studies on respiratory infections.

The analysis found the greatest number of severe cases, requiring hospitalisation, in people in their 50s (222 out of 790 cases), but once

the researchers had adjusted for the fact that many milder cases will have gone undiagnosed, the hospitalisation rate is 8.2%, compared with an estimated 18.4% in the most at risk group, the over 80s (51 out of 76 cases before adjustment). 154 out of 743 people in their 40s had severe symptoms, whereas 133 out of 263 people in their 70s had severe symptoms, but the adjusted hospitalisation rates were again even wider apart than the bare numbers suggest: 4.3% for 40 to 49-year-olds compared with 16.6% for 70 to 79-year-olds. Of those in their 60s, 201 out of 560 cases were severe, whereas the adjusted hospitalisation rate was 11.8%.

The hospitalisation rates were lower for younger age groups: 3.4% of people in their 30s are likely to be hospitalised (while the unadjusted number of severe cases was 124 out of 733 cases), whereas for people in their 20s the rate is likely to be 1.0% (49 out of 437 cases before adjustment). There was only one severe case out of 50 for those aged 10 to 19 and the hospitalisation rate is estimated at 0.04%, whereas none of the 13 cases analysed in the under-10s were severe.

The average time between the first recorded symptoms and death from COVID-19 was estimated to be 17.8 days. The authors note that as the data are from early in the epidemic, more people might die following a longer time lag. Recovering from the disease is estimated to take slightly longer, with patients being discharged from hospital after an average of 22.6 days.

Most people will recover, even from [severe symptoms](#). Death rates from confirmed cases were estimated at 1.38% across all age groups (1,023 out of 44,672 cases in mainland China, with unreported severe cases likely to add to the total, requiring the ratio to be adjusted), but the estimates rise rapidly with age. For example, there were no deaths out of 416 confirmed cases in the under 10-year olds, whereas 13.4% of people aged 80 or above were estimated to die (208 out of 1,408 cases before

adjustment).

The proportion of all people infected who die from the disease—most of whom will display only mild to moderate symptoms—is estimated to be slightly lower, at 0.66%. Again, the risk of [death](#) is much higher in older age groups. For example, 0.031% of people in their 20s are estimated to die, compared with 7.8% of the over-80s.

The authors note that they are unable to adjust for the effect on prognosis of underlying health conditions until individual-level data become available. However, underlying conditions are likely to be correlated with age. Their existence will also vary geographically, particularly between [low-income](#) and high-income regions and countries.

Writing in a linked Comment, Dr. Shigui Ruan (who was not involved in the study) from the University of Miami, USA, says: "Estimates of case fatality ratios might vary slightly from country to country because of differences in prevention, control, and mitigation policies implemented, and because the case fatality ratio is substantially affected by the preparedness and availability of health care. Early studies have shown that delaying the detection of infected cases not only increases the probability of spreading the virus to others (most likely family members, colleagues, and friends) but also makes the infection worse in some cases, thereby increasing the case fatality ratio."

More information: Robert Verity et al, Estimates of the severity of coronavirus disease 2019: a model-based analysis, *The Lancet Infectious Diseases* (2020). [DOI: 10.1016/S1473-3099\(20\)30243-7](https://doi.org/10.1016/S1473-3099(20)30243-7)

Provided by Lancet

Citation: Comprehensive COVID-19 hospitalisation and death rate estimates help countries best prepare as global pandemic unfolds (2020, March 31) retrieved 10 May 2024 from <https://medicalxpress.com/news/2020-03-comprehensive-covid-hospitalisation-death-countries.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.