

Coronavirus: How big, how bad, and what to look out for

March 10 2020, by Ina Skosana



Credit: AI-generated image ([disclaimer](#))

Cases of illness from the new coronavirus (SARS-CoV-2) disease, [known as COVID-19](#), have been confirmed in more than 100 countries. The outbreak was first recorded in China in December last year. [Coronaviruses](#) are a family of viruses that cause illnesses that can range from a common cold to more severe diseases such as Middle East

respiratory syndrome (MERS) and severe acute respiratory syndrome (SARS). The Conversation Africa's Ina Skosana spoke to Shabir Madhi about the situation.

What percentage of people who get the virus are dying from it? Is this high compared with other infectious diseases?

There's no straightforward answer to this question because the epidemic is still at an early stage. It all depends on what the denominator is, which influences the case [fatality](#) risk. And we're still working with unknowns.

The current [estimate](#) is that between 1% and 3% of people diagnosed with COVID-19 die. The problem with this estimate is that the starting point is when individuals have become ill, rather than when they were infected. The actual number of people who are infected might be much higher than the actual number of cases that are presenting for [medical care](#).

The implications of this are huge. For example, if the number of people who have been infected but haven't gone on to develop the disease is high it would mean that the case fatality risk would go down.

The numbers in Iran point to further difficulties with calculating the fatality rate. Initial reports from Iran indicated a case fatality of 10%. This seems very unlikely. This indicates that a large number of cases were going undetected and possibly that mainly very severe cases were being investigated. When investigating very severe cases the case fatality risk is going to be higher.

A 1% fatality risk is moderate and is in the ballpark of other viruses such as respiratory syncytial virus associated with hospitalisation in children.

It is, however, lower than the case fatality risk was for the [coronavirus](#) strains that caused the [SARS](#) epidemic in 2002—which was 10% – and for [MERS in 2013](#), which was about 35%.

Nevertheless, based on current knowledge, the case fatality risk for COVID-19 is higher than observed for seasonal influenza virus, which has a fatality risk of about [0.1%](#). Annually, seasonal influenza virus is estimated to cause [290,000 to 650,000](#) deaths globally. Currently, the global number of deaths due to [COVID-19](#) is more than 3,100.

Is the reaction overblown?

No. It's warranted. This is because we don't know how this epidemic is going to pan out. So even if the case fatality risk is very low, if a high percentage of a population becomes infected that low fatality risk could result in a large number of deaths. For example, if an epidemic involves 100 people with a case fatality risk of 10%, 10 people will die. But if an epidemic infects a million people with a case fatality risk of 1%, there will be 10,000 people dying.

The case fatality risk needs to be interpreted in the context of the number of individuals who are going to become infected to be able to understand the total impact of the epidemic. And right now we don't know what percentage of the different populations will become infected.

Who is most vulnerable to dying from infection?

Current experience—which is largely driven by what has come out of China and more recently from Iran and Italy—is that people over the age of 50 seem to be more susceptible, particularly if they have underlying medical conditions or co-morbid conditions. These include cardiac problems, diabetes and lung disease.

The big unknown for Africa, particularly countries like South Africa, is what the impact will be on populations with a high prevalence of HIV and TB.

The epidemic has only broken out in countries with very low HIV prevalence, so we don't know to what extent individuals living with HIV might have an increased susceptibility to severe disease. This is true of TB too. Because TB affects people's lungs it means that people with the disease have a low tolerance threshold for an additional assault.

In the context of HIV, we have [shown](#) that even in the era of antiretroviral treatment, individuals with HIV still have a 10-fold greater susceptibility to severe influenza illness than the general population, and a higher case fatality risk. We expect there may be differences in susceptibility and outcomes for COVID-19 cases in settings such as South Africa.

What are the symptoms that should get me worried?

The symptoms are very non-specific and very similar to other viral infections like the influenza virus. Unfortunately there isn't a particular trigger that's different to any of the other viruses.

But people can be mindful of a few things. For example, their travel history to a place where the virus is circulating. Or whether the person has come into contact with someone who has visited one of the places where the [virus](#) has been shown to be circulating.

Of course this isn't definitive.

The classical signs and symptoms one can look out for include a fever, cough, headache and myalgia (muscle pain). Again these are very non-descript.

What treatment is available?

Currently there's no specific antiviral treatment to cure SARS-CoV-2 infection. Besides supportive care, no therapeutic agent has been proven to be effective in treating or improving the outcome of COVID-19 cases. There are [clinical trials](#) under way and others are being planned to evaluate different antiviral agents and other possible therapeutic options. The biggest concerns are about a more severe disease, where patients may end up with pulmonary damage which requires supportive care including supplemental oxygen and then might have to be put on to ventilator. Current estimates are that 15% of COVID-19 cases result in severe disease.

Most low- to middle-income settings have fragile health care systems that lack resilience against external shocks such as the demands posed by unexpected epidemics. These countries may lack the capacity to scale up curative services over a relatively short period of time. Such a scale-up of curative services was undertaken in Wuhan to complement existing curative services.

But a quick-fix solution to address an increase in demand for curative health-care services is unlikely in most African countries should SARS-CoV-2 start circulating widely on the continent. This is a culmination of the systematic lack of investment in health care in most countries. It's not something that can be rapidly fixed in the middle of a pandemic.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Coronavirus: How big, how bad, and what to look out for (2020, March 10) retrieved 2

May 2024 from <https://medicalxpress.com/news/2020-03-coronavirus-big-bad.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.