


High-resolution PET/CT assesses brain stem function in patients with hearing impairment

March 25 2020



Axial ^{18}F -FDG PET scans at level of PAC (top, VOI outline in blue) and IC (bottom) in 2 representative patients with left-sided (left image) and right-sided (right image) AHL. Arrows indicate side of decreased metabolism. Credit: Society of Nuclear Medicine and Molecular Imaging

Novel, fully digital, high-resolution positron emission tomography/computed tomography (PET/CT) imaging of small brain stem nuclei can provide clinicians with valuable information concerning the auditory pathway in patients with hearing impairment, according to a new study published in the March issue of *The Journal of Nuclear Medicine*. Using 18F-FDG PET/CT imaging, researchers found that patients with asymmetrical hearing loss have reduced glucose metabolism in parts of the brain stem and primary auditory cortex. The latter may be influenced by cortical reorganization and thus, hopefully help to predict the chance that a cochlear implant will improve hearing.

"With the possible exception of few dedicated high-resolution research scanners, earlier PET/CT systems with lower resolution did not permit clear-cut identification and assessment of [brain](#) stem nuclei," said Iva Speck, MD, resident of otorhinolaryngology at the University of Freiburg Medical Center in Freiburg, Germany. "Today, the use of fully digital clinical PET/CT systems permits greatly enhanced imaging and quantitative assessment of small brain stem nuclei, such as the inferior colliculus (IC), the part of the midbrain that acts as a main auditory pathway for the body."

In the study, 13 patients with asymmetric [hearing](#) loss underwent 18F-FDG PET/CT imaging. The scans were reviewed by two experienced readers who examined regional [glucose metabolism](#) in the IC and the [primary auditory cortex](#) (PAC)—a part of the brain known to undergo metabolic changes based on acoustical outside input and transformation to neuronal signals from the cochlea hair cells to the auditory nerve fibers. The readers rated the scans as to whether glucose [metabolism](#) showed no asymmetry or mild, moderate or strong asymmetry to the left or to the right for the IC and PAC separately. Statistical analyses were performed to determine the effect of the duration of [hearing impairment](#) on glucose metabolism and to compare glucose metabolism between the IC and PAC.

Regional glucose metabolism of both the IC and PAC was significantly reduced on the contralateral (opposite) side of the poorer-hearing ear, as compared to the ipsilateral (same) side. In addition, a longer duration of hearing impairment was associated with a higher metabolism on the contralateral PAC. By contrast, duration of hearing impairment did not predict regional glucose metabolism for the ipsilateral PAC or either side of the IC.

"Previous studies suggest that the association between longer duration of hearing impairment and higher [glucose](#) metabolism indicates cortical reorganization. In bilateral deaf patients this has been shown to lessen the benefits of cochlear implants," said Speck. "Prediction of a successful cochlear implant outcome might benefit from improved imaging with fully digital PET/CT systems, as large parts of the auditory system, including small brain nuclei such as the IC, can be assessed for preoperative patient characterization."

She continued, "Beyond this topic, the study's findings are of interest for other neurological research fields, like neurodegenerative diseases, which often affect brain stem nuclei early in disease course," Speck remarked. "Digital PET pushes the limits of what can be imaged and contributed to patient care by molecular imaging."

More information: Iva Speck et al, 18F-FDG PET Imaging of the Inferior Colliculus in Asymmetric Hearing Loss, *Journal of Nuclear Medicine* (2019). [DOI: 10.2967/jnumed.119.231407](https://doi.org/10.2967/jnumed.119.231407)

Provided by Society of Nuclear Medicine and Molecular Imaging

Citation: High-resolution PET/CT assesses brain stem function in patients with hearing impairment (2020, March 25) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2020-03-high-resolution-petct-brain-stem-function.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.