


# Homeless people receive less treatment in hospitals for heart attacks, have higher readmission rates

March 19 2020

---



Credit: CC0 Public Domain

Homelessness has become a social crisis and public health problem around the world, affecting people of all ages. Most homeless people are at a disadvantage with few resources, and may or may not have adequate

health insurance. Mental illness and substance abuse are common issues in the homeless community. People living on the streets have a high likelihood of developing heart disease. Yet they have little ability to take care of their health.

A new study published in *Mayo Clinic Proceedings* examined in-[hospital outcomes](#) and 30-day readmission rates of 3,937 [homeless patients](#) from more than 1.5 million people admitted for a heart attack. Researchers used two years of data (2015-2016) from the National Readmission Database. The findings show that homeless patients often receive less treatment and are more likely to end up readmitted to the hospital within a month.

"The differences in patterns of care and clinical outcomes between homeless and nonhomeless patients are likely a reflection of the differences in risk profile at baseline," says Mohamad Alkhouli, M.D., a Mayo Clinic cardiologist and senior author of the study.

Compared to those with stable housing, homeless patients admitted with a heart attack tended to be male, about 10 years younger, and had fewer traditional risk factors for [heart disease](#). Homeless patients in the study had a higher incidence of congestive heart failure, anemia, chronic kidney disease, and liver and lung diseases. Anxiety, depression, [substance abuse](#) and HIV infections were substantially higher for this group, as well.

Looking at matched outcomes for each group, homeless patients received less treatment in the hospital, as shown through a lower rate of angiography and revascularization. The in-hospital mortality rate and median cost of hospitalization were similar for the homeless versus patients who were not homeless.

Other differences emerged when comparing length of stay and

readmission. Homeless patients were in the hospital longer. Yet they were more likely to leave against medical advice and be discharged to an intermediate care facility. They were also more likely to be readmitted. The 30-day readmission rates differed significantly, with 22.5% of homeless patients readmitted, compared to only 10% of patients who were not homeless. Cardiovascular causes were the most common reasons for readmission in both groups. However, 18% of homeless patients were readmitted for psychiatric causes, compared to 2% of patients who were not homeless.

"Addressing cardiovascular disease in the growing population of the homeless comes with a unique set of challenges," says Dr. Alkhouli. "The high burden of psychiatric illness among the homeless emphasizes the need for a [holistic approach](#) to address both the clinical and social needs of these patients."

**More information:** R. Scott Wright et al. Homeless With a Heart Attack in America—A Double Whammy, *Mayo Clinic Proceedings* (2020). [DOI: 10.1016/j.mayocp.2020.02.022](https://doi.org/10.1016/j.mayocp.2020.02.022)

Provided by Mayo Clinic

Citation: Homeless people receive less treatment in hospitals for heart attacks, have higher readmission rates (2020, March 19) retrieved 3 May 2024 from <https://medicalxpress.com/news/2020-03-homeless-people-treatment-hospitals-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---