

Getting too little—or too much—sleep may be bad for the heart

March 18 2020



Credit: CC0 Public Domain

Whether you like to burn the midnight oil to check emails or binge watch your favorite series, toss and turn or sleep until mid-morning, it seems the amount of sleep you get matters when it comes to your future

vascular and heart health. Compared with people who slept for longer or shorter periods of time, those who reported sleeping seven or eight hours a night had significantly less evidence of stiffness in their arteries, indicating a lower chance of developing heart disease or suffering a stroke, according to research presented at the American College of Cardiology's Annual Scientific Session Together with World Congress of Cardiology (ACC.20/WCC).

Even after accounting for other known [risk factors](#) for heart disease or stroke, people who slept less than six hours or more than eight hours a night had significantly greater odds of having plaque buildup in the walls of their carotid arteries—a 54% and 39% increase, respectively—compared with those who got seven or eight hours of shut eye. The study adds to mounting evidence that [sleep patterns](#), similar to diet and exercise, may play a defining role in someone's cardiovascular risk.

"The message, based on our findings, is 'sleep well, but not too well.' Getting too [little sleep](#) appears bad for your health but too much seems to be harmful as well," said Evangelos Oikonomou, MD, consultant cardiologist and the study's lead author. "Unlike other heart disease risk factors such as age or genetics, [sleep habits](#) can be adjusted, and even after taking into consideration the impact of established risk factors for atherosclerosis and cardiovascular diseases—for example age, gender, obesity, smoking, hypertension, diabetes, [high blood pressure](#) and even a history of coronary artery disease—both short and long sleeping [duration](#) may act as additional risk factors."

For this analysis, researchers assessed sleep patterns in 1,752 people living in the Corinthia region of Greece using a standard questionnaire that was fielded by a trained cardiologist, primary care provider or nurse. Participants were then divided into one of four groups based on self-reported [sleep duration](#): normal (seven to eight hours a night), short sleep

duration (six to seven hours a night), very short sleep duration (less than six hours a night) or long sleep duration (greater than eight hours a night). Participants represented a broad spectrum of the general public, including healthy people as well as those with cardiovascular risk factors and established heart disease, and most were from rural areas with less than 1,000-2,000 inhabitants. They ranged in age from 40 to 98 years, with a mean age of 64 years old.

At the time of the study, each participant also underwent ultrasound imaging to measure the thickness of the inner part of the arterial wall. Thickening of the arterial walls reflects plaque buildup and is associated with an increased risk of stroke and other cardiovascular events. Intima media thickness of >1.5 mm or protrusion $>50\%$ compared to nearby segments of the artery wall was defined as atherosclerotic plaque.

Researchers uncovered a U-shaped pattern between sleep duration and early indicators of atherosclerosis, which underscores the need for a balanced sleep pattern, Oikonomou said. Intima media thickness and plaque build-up in the artery walls was greater in both the shorter and longer sleep duration groups as compared to normal sleep duration.

"We don't fully understand the relationship between sleep and cardiovascular health. It could be that sympathetic nervous system withdrawal or a slowing [of this system] that occurs during sleep may act as a recovery phase for [usual] vascular and cardiac strain," Oikonomou said. "Moreover, short sleep duration may be associated with increased cardiovascular risk factors—for example, unhealthy diet, stress, being overweight or greater alcohol consumption—whereas longer sleep duration may be associated with a less active lifestyle pattern and lower physical activity."

Researchers said that adopting a balanced sleep pattern of six to eight hours nightly may be just what the doctor ordered.

"It seems that this [amount of sleep](#) may act as an additive cardioprotective factor among people living in modern western societies, and there can be other health benefits to getting sufficient and quality sleep," Oikonomou added.

The amount of sleep someone needs depends on several factors, especially one's age. In the U.S., most guidelines recommend that adults sleep between seven and nine hours each night. Yet, one out of every three American adults do not get enough sleep, according to the Centers for Disease Control and Prevention. Poor sleep has also been linked to a higher risk of obesity, diabetes, high blood pressure, [heart disease](#), poor mental health and even dying early.

This study is limited in that it relies on self-reported sleep patterns and is cross sectional in nature, so the relationship between sleep patterns and atherosclerotic activity is based on a single point in time. Further research is needed, especially to look at whether too much [sleep](#) is harmful, which hasn't been studied as well as getting too little.

Provided by American College of Cardiology

Citation: Getting too little—or too much—sleep may be bad for the heart (2020, March 18) retrieved 5 May 2024 from

<https://medicalxpress.com/news/2020-03-littleor-muchsleep-bad-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--