

# Researchers identify protein critical for wound healing after spinal cord injury

March 2 2020

---



Credit: CC0 Public Domain

Plexin-B2, an axon guidance protein in the central nervous system (CNS), plays an important role in wound healing and neural repair following spinal cord injury (SCI), according to research conducted at

the Icahn School of Medicine at Mount Sinai and published today in *Nature Neuroscience*. The study's findings could aid the development of therapies that target axon guidance pathways for more effective treatment of SCI patients.

Tissue repair after SCI requires the mobilization of immune and [glial cells](#) to form a protective barrier that seals the wound, facilitates debris clearing and contains inflammation. Building this barrier involves a process called corralling wherein microglia ([immune cells](#) in the CNS) and macrophages (immune cells that originate from blood) form a barrier around the lesion that separates healthy and necrotic tissue. In this study, researchers found that this corralling begins early in the healing process and requires Plexin-B2, a protein that facilitates the movement of immune cells by steering them away from colliding cells.

Researchers found that the deletion of Plexin-B2 in microglia and macrophages impaired corralling, which led to [tissue damage](#), inflammatory spillover, and hindered the regeneration of axons (slender part of a nerve cell where impulses are conducted).

"The role of microglia and macrophages in the spatial organization of glial cells around the injury site via an axon guidance receptor is quite unexpected" said lead investigator Hongyan Jenny Zou, MD, Ph.D., Professor of Neurosurgery and Neuroscience at the Icahn School of Medicine at Mount Sinai.

Tissue repair in the CNS relies on a coordinated response from diverse cell types in overlapping phases. This complexity makes it difficult to distinguish specific roles of glial cell populations. Previously, astrocytes (supporting glial cells) were presumed to be the main driver for corralling. However, this study identified the critical contribution of injury-activated microglia/macrophages, as well as the role of Plexin-B2 in corralling. Understanding the signaling pathways and interactions of

glial cells with each other and the injury environment is fundamental to improving [neural repair](#) after a traumatic brain or spinal cord injury.

**More information:** Xiang Zhou et al. Microglia and macrophages promote corralling, wound compaction and recovery after spinal cord injury via Plexin-B2, *Nature Neuroscience* (2020). [DOI: 10.1038/s41593-020-0597-7](#)

Provided by The Mount Sinai Hospital

Citation: Researchers identify protein critical for wound healing after spinal cord injury (2020, March 2) retrieved 6 May 2024 from <https://medicalxpress.com/news/2020-03-protein-critical-wound-spinal-cord.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--