


# The origin of satiety: Brain cells that change shape after a meal

March 3 2020



Remodelling of the satiety circuit of POMC neurons after a balanced meal. Red box: area corresponding to the photo at right. Credit: Alexandre Benani / CNRS / CSGA


Researchers from the CNRS, Inrae, University of Burgundy, Université de Paris, Inserm, and University of Luxembourg have just revealed the mechanisms in the brain that lead to feelings of satiety after eating. They involve a series of reactions triggered by a rise in blood glucose levels. This study, which was conducted on mice, is published in *Cell Reports* on 3 March 2020.

The [neuronal circuits](#) in the [brain](#) governing feelings of hunger and satiety can modify their connections, thereby adjusting feeding behavior to living conditions and maintaining a balance between food intake and calorie expenditure. Scientists suspect that this plasticity could be altered for obese subjects.

In a new study conducted on mice, a team led by Alexandre Benani, a CNRS researcher at the Centre for Taste and Feeding Behavior (CNRS/Inrae/University of Burgundy/AgroSup Dijon), has shown that these circuits are activated on the time scale of a meal, subsequently regulating feeding behavior. However, this activation does not occur through a change in the circuit's "connections."

Scientists focused on POMC neurons in the hypothalamus, located at the base of the brain, which are known for limiting food intake. They are connected to a large number of neurons from other parts of the brain, with the connections of this circuit being malleable: They can be made and unmade very quickly based on hormonal fluctuations. Researchers observed that this neuronal circuit is not modified after a balanced meal, but that other [nerve cells](#) associated with POMC neurons, known as

astrocytes, actually change form.



POMC neurons (orange dots) in the hypothalamus of a mouse, located at the base of the brain. Photo taken from a mouse using a confocal microscope. Credit: Danaé Nuzzaci / CNRS / CSGA

Astrocytes are star-shaped nerve cells that were first studied for their supporting role with respect to neurons. Under usual conditions, they sheathe POMC neurons and act somewhat like [brake pads](#) by limiting their activity. After a meal, [blood glucose levels](#) (glycemia) temporarily increase, with astrocytes detecting this signal and retracting in less than

one hour: once this "brake" is released, POMC [neurons](#) are activated, ultimately promoting the feeling of satiety.

Surprisingly, a meal that is high in fats does not lead to this remodeling. Does this mean that lipids are less effective in satisfying hunger? The scientists are trying to determine whether they trigger satiety through another circuit. It also remains to be seen whether sweeteners have the same effects, or whether they lure the brain by providing an addictive sensation of sweetness without satisfying hunger.

**More information:** *Cell Reports* (2020). [DOI: 10.1016/j.celrep.2020.02.029](#)

Provided by CNRS

Citation: The origin of satiety: Brain cells that change shape after a meal (2020, March 3)  
retrieved 7 May 2024 from  
<https://medicalxpress.com/news/2020-03-satiety-brain-cells-meal.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------