


Brain insulin sensitivity determines body weight and fat distribution

April 24 2020



If the brain reacts sensitively to the hormone, you lose a significant amount of weight, reduce unhealthy abdominal fat and can maintain weight in the long term. If the brain reacts only slightly or not at all to insulin, you only lose some weight at the beginning of the intervention and then gain weight again. Credit: IDM

Just where fat is deposited in the body and to what degree a person may benefit from a lifestyle intervention depends, among other things, on how sensitive the brain is to insulin. If the person's brain responds sensitively to the hormone, a significant amount of weight can be lost, unhealthy visceral fat reduced, and the weight loss can be maintained over the long term. However, If the person's brain responds only slightly or not at all to insulin, the person only loses some weight at the beginning of the intervention and then experiences weight regain. Over the long term, visceral fat also increases. These are the results of a long-term study by the German Center for Diabetes Research (DZD), Helmholtz Zentrum München and Tübingen University Hospital which has now been published in *Nature Communications*.

To what extent body fat has an unhealthy effect depends primarily on where it is stored. If fat accumulates in the abdomen, this is particularly unfavorable. This is because the visceral fat releases numerous neurotransmitters that affect blood pressure, influence the secretion of the hormone insulin and can cause inflammation. This increases the risk of diabetes, cardiovascular disease and certain types of cancer. The subcutaneous fat which accumulates on the buttocks, thighs and hips has no adverse health effects. However, it is still unclear why fat storage does not occur in the same place in all people.

Studies in the Tübingen Lifestyle Intervention Program (TULIP) suggest that brain insulin responsiveness could play an important role here. They showed that people with a high insulin [sensitivity](#) in the brain benefit significantly more from a [lifestyle intervention](#) with a diet rich in fiber and exercise than people with [insulin resistance](#) in the brain. Not only did they lose more weight, they also had a healthier fat distribution. But how does insulin sensitivity affect the distribution of body fat and weight in the long term?

Researchers from the German Center for Diabetes Research (DZD),

Helmholtz Zentrum München and Tübingen University Hospital investigated this question in a long-term study. For this purpose, they recorded the follow-up data of 15 participants over a period of nine years, in which the insulin sensitivity in the brain was determined by magnetoencephalography before the start of a 24-month lifestyle intervention.

High insulin sensitivity associated with reduction in visceral fat and weight

It was found that insulin action in the brain not only determines body weight, but also the distribution of fat in the body. "Subjects with high insulin sensitivity in the brain benefited from the lifestyle intervention with a pronounced reduction in weight and visceral fat. Even after the lifestyle intervention had ended, they only regained a small amount of fat during the nine-year follow-up," said the head of the study, Professor Martin Heni from Tübingen University Hospital. In contrast, people with brain insulin resistance only showed a slight [weight loss](#) in the first nine months of the program. "Afterwards, their body [weight](#) and visceral fat increased again during the following months of lifestyle intervention," said first author PD Dr. Stephanie Kullmann from the IDM.

Since insulin action in the hypothalamus is crucial for the regulation of peripheral energy metabolism, the researchers also investigated how insulin sensitivity in this area of the brain is related to the distribution of body fat. For this purpose, they examined a cross-sectional cohort of 112 participants. The analysis of the data showed that people with high insulin sensitivity in the hypothalamus form little visceral fat. However, [insulin sensitivity](#) has no influence on the mass of subcutaneous fat.

"Our study reveals a novel key mechanism that regulates fat distribution in humans. Insulin sensitivity in the brain determines where fat is

deposited," said Heni, summarizing the results. Since [visceral fat](#) not only plays a role in the development of type 2 diabetes, but also increases the risk of cardiovascular disease and cancer, the study results may also open up new approaches for treatment options beyond metabolic diseases. The researchers in Tübingen are already working on new therapies to abolish [insulin](#) resistance in the [brain](#) and thus have a beneficial effect on [body](#) fat distribution.

More information: Stephanie Kullmann et al, Brain insulin sensitivity is linked to adiposity and body fat distribution, *Nature Communications* (2020). [DOI: 10.1038/s41467-020-15686-y](https://doi.org/10.1038/s41467-020-15686-y)

Provided by Deutsches Zentrum fuer Diabetesforschung DZD

Citation: Brain insulin sensitivity determines body weight and fat distribution (2020, April 24) retrieved 6 May 2024 from <https://medicalxpress.com/news/2020-04-brain-insulin-sensitivity-body-weight.html>

| |
|---|
| This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only. |
|---|