

Cardiorespiratory fitness assessment improves accuracy of health predictions

April 30 2020



Credit: CC0 Public Domain

According to a new study in *Mayo Clinic Proceedings*, taking cardiorespiratory fitness (CRF) into account along with traditional risk factors such as age, sex, blood pressure, cholesterol, and smoking, improves the accuracy of mortality risk assessment.



"We found that within a contemporary adult UK population with a varied and relatively low pre-test risk profile, greater CRF was strongly associated with a lower risk of mortality. The findings enabled new insights into the incremental prognostic value of a CRF evaluation," explained lead investigator Jari A. Laukkanen, MD, Ph.D., Cardiologist, Institute of Public Health and Clinical Nutrition, University of Eastern Finland, Kuopio, Finland. "The predictive accuracy of CRF levels was even more evident when factored in with traditional <u>risk factors</u> such as age, sex, <u>blood pressure</u>, cholesterol and smoking."

Although conventional risk factors such as age, sex, blood pressure, cholesterol, and smoking can significantly contribute to the risk of vascular disease and mortality, these established risk factors are often absent in a large proportion of cardiovascular disease patients. This contributes to difficulties in identifying individuals at increased risk for vascular disease and/or death.

CRF is recognized as an important marker of both functional ability and mortality and a wealth of literature shows this measure to be inversely and independently associated with vascular disease and mortality. However, physicians do not routinely assess fitness—in either general or specialized clinical settings—when testing for traditional risk factors.

This study was based on data from the large-scale UK Biobank Study, which gathered information on close to 60,000 participants aged 40-69 years. It assessed CRF using submaximal exercise testing in the form of a six-minute stationary bicycle protocol. This protocol is easily available, has a good safety profile, reliability, and validity. The investigators evaluated the impact of the CRF assessment against and in conjunction with risk factors typically measured in clinical practice. Investigators used mortality data from the National Health Service Information Centre (England and Wales) and the Scottish Morbidity Record, from which they were able to classify the causes of death.



This is one of the first large scale population-based studies showing that risk prediction can be improved in a pre-screened relatively low risk population by adding information on CRF, estimated on the basis of a submaximal exercise test, to conventional cardiovascular risk factors. During a median follow-up of 5.8 years, 936 deaths occurred within the study group.

"Do we really need to make CRF evaluations a standard part of a medical exam? Based on what we learned, the answer is 'yes.' Our findings clearly validate them as vital clinical risk predictors in conjunction with the other factors," noted Dr. Laukkanen. "Submaximal exercise testing in order to factor in CRF levels with conventional risk factors should be routinely used by clinicians to determine whether highrisk patients need additional interventions."

In an accompanying <u>editorial</u>, Carl J. Lavie, MD, Cardiologist, John Ochsner Heart and Vascular Institute, Ochsner Clinical School—University of Queensland School of Medicine, New Orleans, LA, USA, and co-authors observed that "Certainly, numerous studies show the powerful impact of CRF in predicting risk. Although the gold standard assessment of CRF is cardiopulmonary exercise testing and using gas exchange to measure peak oxygen consumption, many clinicians do not have easy access to such testing, and neither do they have the equipment and personnel to monitor these tests, not to mention the moderate expense of this testing. The unique aspect of the present UK Biobank study was using a submaximal six-minute bicycle test in a very large cohort to assess CRF and mortality combined with established CVD risk factors."

CRF is an index of habitual physical activity, particularly aerobic exercise. Good CRF helps to ward off chronic diseases, provides general wellbeing, and reduces the risk of premature death. In addition to good nutrition and, adequate rest and relaxation, individuals can improve their



fitness levels by undertaking regular exercise. Fitness and exercise are potential ways to avoid daily stress, which can take a substantial toll on human health.

"We need to do a better job promoting physical activity and exercise training (aerobic and resistance) throughout the healthcare system, schools, the workplace, and society for the primary and secondary prevention of cardiovascular disease. The long-term health of our populations will depend on the success of these efforts," commented Dr. Lavie and co-authors.

More information: "Prognostic Relevance of Cardiorespiratory Fitness as Assessed by Submaximal Exercise Testing for All-Cause Mortality: A UK Biobank Prospective Study,", *Mayo Clinic Proceedings* (2020). DOI: 10.1016/j.mayocp.2019.12.030

Provided by Elsevier

Citation: Cardiorespiratory fitness assessment improves accuracy of health predictions (2020, April 30) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2020-04-cardiorespiratory-accuracy-health.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.