


Reducing early brain inflammation could slow Alzheimer's progression

April 27 2020



This fluorescent photomicrograph shows astrocytes (green) and neurons (pink) in the dentate gyrus of rat hippocampus. Credit: Giorgia Menegoni, Sapienza, University of Rome

In a new animal study examining Alzheimer's disease, researchers found that disease progression could be slowed by decreasing neuroinflammation in the brain before memory problems and cognitive impairment were apparent. The new findings point to the importance of developing therapies that target very early stages of the disease.

In 2011, the National Institute on Aging updated the [diagnostic criteria](#) for Alzheimer's [disease](#) to reflect its progressive nature. The criteria added a preclinical stage during which brain changes are taking place, but the person is still asymptomatic and, therefore, unaware of his condition. Biomarker profiles could eventually be used to identify people in the disease's early stages who might benefit from early treatments.

"Starting an intervention at the earliest stage of the disease, when cellular and molecular alterations have already been triggered but major damage to the brain has not yet occurred, could offer a way to reduce the number of people who go on to develop full Alzheimer's dementia," said research team leader Caterina Scuderi, Ph.D., assistant professor of pharmacology and toxicology from Sapienza, University of Rome. "However, there have been few studies in [animals](#) examining therapeutic strategies that target timepoints before symptoms can be seen."

Scuderi was scheduled to present this research at the American Association for Anatomy [annual meeting](#) in San Diego this month. Though the meeting, to be held in conjunction with the 2020 Experimental Biology conference, was canceled in response to the COVID-19 outbreak, the research team's abstract was published in this month's issue of *The FASEB Journal*.

The researchers designed an animal study to gain a deeper understanding of [neuroinflammation](#)'s role in Alzheimer's disease during the pre-symptomatic stage of the disease, which might represent the best time

for therapeutic intervention. The study results suggest that rebalancing neuroinflammation in animals that show altered neuroinflammatory parameters could be beneficial.

"Our results help demonstrate that neuroinflammation in Alzheimer's disease is an extremely complex phenomenon that can change over the disease's progression and varies based on factors such as affected brain area," said Scuderi. "We hope that these findings will prompt scientists to further investigate neuroinflammation at the earliest stages of the disease, which may represent an important pharmacological target."

Provided by Experimental Biology

Citation: Reducing early brain inflammation could slow Alzheimer's progression (2020, April 27) retrieved 5 May 2024 from

<https://medicalxpress.com/news/2020-04-early-brain-inflammation-alzheimer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--