

Study shows ICU patients with low-risk penicillin allergies can be tested and treated

April 3 2020



Lead author Cosby A. Stone, Jr., MD, instructor in Allergy/Immunology at Vanderbilt University Medical Center. Credit: Vanderbilt University Medical Center

Many patients previously diagnosed with a penicillin allergy can have their allergy label removed after testing and safely undergo treatment with penicillin medications, according to a study published in *American*

Journal of Respiratory and Critical Care Medicine.

Around 8-15% of the U.S. population is labeled with a [penicillin allergy](#), but many of these allergies are the result of viral rashes, drug-viral interactions, or non-allergic side effects. Even [patients](#) who experience a severe anaphylactic reaction appear to lose their sensitivity at a rate of 10% or more every year.

"To date, our team has removed more than 90 low-risk penicillin allergies without any patients reporting a symptomatic challenge," said lead author Cosby A. Stone, Jr., MD, instructor in Allergy/Immunology at Vanderbilt University Medical Center.

"Around one-third of the patients whose penicillin allergies were removed have already gone on to safely use penicillin treatments in their subsequent health care when they were needed," he said.

A false penicillin label can prohibit appropriate patient treatment through limitations that force doctors to use broader spectrum and second line antibiotics, increasing the chances of surgical site infections, greater health care utilization, treatment failure for common infections, drug resistant infections, and longer lengths of stay.

Data collected from the outpatient Drug Allergy Clinic at Vanderbilt University Medical Center was used to develop a risk-stratification tool that could identify patients with low-risk penicillin allergies in the Medical Intensive Care Unit (MICU).

Over a seven-month period, patients who were screened and identified as low-risk received an oral dose of 250mg of amoxicillin and remained under observation for an hour and a half following this challenge, without a preceding allergy skin test.

The team's results showed that none of the low-risk patients who underwent a direct oral challenge during the study period had any symptoms of an allergic reaction when challenged, enabling their [penicillin allergy](#) label to be removed.

Directly challenging low-risk penicillin allergies with a dose of amoxicillin is expected to be a strategy that can remove up to 60% of all penicillin allergies and improve a patient's future health care by expanding possible treatment options.

More information: Cosby A Stone et al, Risk-Stratified Management to Remove Low-Risk Penicillin Allergy Labels in the Intensive Care Unit, *American Journal of Respiratory and Critical Care Medicine* (2020). [DOI: 10.1164/rccm.202001-0089LE](https://doi.org/10.1164/rccm.202001-0089LE)

Provided by Vanderbilt University Medical Center

Citation: Study shows ICU patients with low-risk penicillin allergies can be tested and treated (2020, April 3) retrieved 9 May 2024 from <https://medicalxpress.com/news/2020-04-icu-patients-low-risk-penicillin-allergies.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.