

Interventions boost abstinence, condom use for black teens

April 20 2020



Credit: CC0 Public Domain

Sexual health interventions are effective at increasing both abstinence and condom use in Black adolescents, according to research from North

Carolina State University evaluating dozens of studies on interventions and outcomes in Black youth.

The new paper, published in *JAMA Pediatrics*, draws on data from 29 studies that reported on a total of 11,918 Black adolescents. Sexual health interventions range from school-based health classes to programs offered by community organizations.

"We focused on Black adolescents because they face greater health disparities when it comes to the risk of unplanned pregnancy and contracting sexually-transmitted infections (STIs) compared to other adolescents," says Reina Evans, first author of the paper and a Ph.D. student at NC State. "This disparity stems, in large part, from the context in which Black teens make decisions about their health. For example, stress from racism and discrimination, as well as unequal access to health care can impact the health of Black teens. We wanted to see whether sexual health interventions can be a [valuable tool](#) in addressing this disparity."

The researchers found that [young people](#) were slightly more likely to abstain from sex if they took part in one of these programs—particularly if the [intervention](#) occurred at school. The researchers also found a modest increase in condom use for adolescents who took part in an intervention.

"When we considered if youth who completed these programs had lower rates of unintended pregnancy or STIs, we did not see any effect—positive or negative—from the programs," Evans says. "However, only 4 of the 29 studies investigated these important outcomes, so these findings should be considered preliminary. It is likely that offering a sexual health program to youth could be one component of the puzzle of preventing teen pregnancy and STIs, and other components are also needed—like increasing access to contraception and

STI testing and addressing racism in healthcare in the United States."

The study also showed that interventions made Black adolescents more knowledgeable about sexual health and more confident that they could and would make safe sexual decisions in the future.

In addition, researchers found that the effects were similar regardless of whether interventions were coeducational or aimed at a specific gender, and regardless of the target age range for an intervention.

"These findings highlight that sexual health interventions make a positive difference for many Black adolescents," Evans says. "Important next steps include finding ways to disseminate interventions more broadly to teens, ensuring the interventions are accessible, and encouraging participation and engagement.

"What's more, there's an urgent need to address disparities in access to long-term, high-quality reproductive healthcare, which could have a significant effect on health outcomes for Black teens."

More information: *JAMA Pediatrics* (2020). [DOI: 10.1001/jamapediatrics.2020.0382](https://doi.org/10.1001/jamapediatrics.2020.0382)

Provided by North Carolina State University

Citation: Interventions boost abstinence, condom use for black teens (2020, April 20) retrieved 18 April 2024 from <https://medicalxpress.com/news/2020-04-interventions-boost-abstinence-condom-black.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.