

# Does long-term exposure to air pollution lead to a steeper rate of cognitive decline?

April 8 2020

---



Credit: CC0 Public Domain

People who live in urban areas with higher levels of air pollution may score lower on thinking and memory tests and may also lose cognitive skills faster over time, or it is possible they also may not, according to a

study published in the April 8, 2020, online issue of *Neurology*, the medical journal of the American Academy of Neurology. Researchers examined the association of air pollution levels and cognitive impairment and decline in participants in two large epidemiological studies. They found an association between the air pollution and cognitive decline in one study group but not in the other.

"As people live longer lives and the [aging population](#) grows, age-related cognitive decline is a growing public health concern with profound social, economic and [health effects](#), so finding ways to reduce the risk is important," said study author Erin R Kulick, Ph.D., M.P.H., of Brown University School of Public Health in Providence, R.I., and a member of the American Academy of Neurology. "Air pollution can affect large populations of people because it has known cardiovascular risks, and previous research has found that it may also contribute to cognitive decline. However, the results of our research were mixed."

The study involved people living in the Northern Manhattan area of New York City who were enrolled in two larger long-term studies: 5,330 people with an average age of 75 enrolled in the Washington Heights-Inwood Community Aging Project; and 1,093 people with an average age of 70 enrolled in the Northern Manhattan Study. Both groups were ethnically and racially diverse with black, white and Hispanic participants.

All participants were given medical exams at the beginning of the study as well as cognitive tests to measure memory, language skills and executive function, which involves thinking skills like organizing, planning and completing tasks.

The group from the Washington-Heights-Inwood study of 5,330 participants was followed an average of seven years with six rounds of follow-up testing every 18 months to two years. The Northern Manhattan

study group of 1,093 participants was followed for five years with one follow-up round of testing.

Researchers used the residential addresses of each participant to determine their exposures to three air pollutants. Those pollutants were [nitrogen dioxide](#) and two groups of particulate matter, particles of liquids or solids suspended in air that were less than 2.5 microns in diameter called fine particulate matter and particles that were less than 10 microns in diameter called respirable particulate matter. Average levels of air pollutants were similar for both groups of participants. In each group, participants were divided into four groups based on their air pollution exposure.

Researchers found that participants in the Washington-Heights Inwood study were exposed to a yearly average of 32 parts per billion of nitrogen dioxide, 13 micrograms per cubic meter ( $\mu\text{g}/\text{m}^3$ ) of fine particulate matter and 21  $\mu\text{g}/\text{m}^3$  of respirable particulate matter. The U.S. Environmental Protection Agency (EPA) considers up to 53 parts per billion to be a safe level of yearly average exposure to nitrogen dioxide, up to 12  $\mu\text{g}/\text{m}^3$  for [fine particulate matter](#) and up to 50  $\mu\text{g}/\text{m}^3$  for respirable particulate matter.

"While the levels were in the range considered safe by the EPA, it's possible that these results reflect higher levels from an earlier point in time. It also raises the question of whether the federal levels are low enough to protect people's health," said Kulick.

The researchers found that in the Washington Heights-Inwood group, people with greater exposure to higher levels of air pollution had lower scores on the tests at the beginning of the study and more rapid rates of decline.

Exposure to nitrogen dioxide was linked to an accelerated rate of

cognitive decline comparable to one year of aging. Results were similar for fine and respirable particulate matter.

In the Northern Manhattan group, researchers did not find an association between cognitive function and air pollution. Kulick says the difference in results may be because the second group was much smaller and only had one round of follow-up compared to six rounds for the first group.

"A strength of our study was that we were able to analyze the rates of [cognitive decline](#) over time so it adds important findings to the growing body of scientific evidence about air pollution and its effects on the brain health of older adults," Kulick said. "More research is needed to better understand our study's mixed results. The good news for public health is that air pollution can be reduced, and has been in some cities, through laws and regulation. But there are still millions of people living in areas of the United States where major air quality improvements are needed."

A limitation of the study was that while it included levels of air pollution near a person's residence, it did not account for levels of pollution at the workplace or elsewhere. Also, many participants grew up in other countries and may have had different exposures to air [pollution](#) at younger ages.

Provided by American Academy of Neurology

Citation: Does long-term exposure to air pollution lead to a steeper rate of cognitive decline? (2020, April 8) retrieved 30 April 2024 from <https://medicalxpress.com/news/2020-04-long-term-exposure-air-pollution-steeper.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--