


Paternal age and the risk of neurodevelopmental disorders in children

April 21 2020



Credit: Tohoku University

Genetic factors play a role in determining whether children have neurodevelopmental disorders. Maternal exposure to drugs and viral or bacterial illnesses can be detrimental, too.

However, a recent epidemiological survey of approximately 6 million people worldwide has revealed that advanced paternal age is associated with the development of [neurodevelopmental disorders](#). In other words,

the older the parent, the [increased risk](#) a child has of developing disorders such as autism, ADHD and other learning disabilities.

A research team from the Department of Developmental Neuroscience at the Tohoku University Graduate School of Medicine has revealed further details about this phenomenon with their recent publication in *PLOS ONE*.

The research team, led by doctoral student Misako Tatehana and Professor Noriko Osumi, performed immunohistochemical analysis of the testis in three-month-old mice before performing the same analysis on mice aged 12 months or older.

Tatehana and her team analyzed histone proteins during the 12-step spermatogenesis process. Histone proteins undergo chemical modifications during spermatogenesis, thereby affecting [gene expression](#). More specifically, the team looked at the seven methylations and one acetylation.

They catalogued these as epigenetic markers—modifications that affect the expression of the gene without changing the DNA base sequence of the genome itself.

Comparisons of the markers between the younger and older mice using imaging quantification techniques revealed that the latter had higher amounts of the modified histone protein, H3K79me3. A previous study by Professor Osumi found a correlation between the amount of H3K79me3 in sperm and abnormal speech communication in pups, making it a predictive marker of neural-developmental disorders.

This new finding signals that paternal aging potentially affects neurodevelopment in humans. Further research on the matter hopes to develop greater diagnostic methods for [disorders](#) stemming from the risk

of advanced paternal age.

More information: Misako Tatehana et al. Comprehensive histochemical profiles of histone modification in male germline cells during meiosis and spermiogenesis: Comparison of young and aged testes in mice, *PLOS ONE* (2020). [DOI: 10.1371/journal.pone.0230930](https://doi.org/10.1371/journal.pone.0230930)

Provided by Tohoku University

Citation: Paternal age and the risk of neurodevelopmental disorders in children (2020, April 21) retrieved 1 May 2024 from <https://medicalxpress.com/news/2020-04-paternal-age-neurodevelopmental-disorders-children.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--