


# Caring for those most vulnerable to COVID-19 requires vigilance

April 16 2020

---



Joseph G. Ouslander, M.D., an internationally renowned geriatrician, a professor of geriatric medicine, and a senior advisor to the dean at FAU's Schmidt College of Medicine. Credit: Florida Atlantic University

As the coronavirus disease (COVID-19) pandemic advances so does the uncertainty and evolution of the pandemic as it relates to nursing homes and long-term care facilities (LTCF). Early reports suggest the case fatality rate for those over 80, which constitutes nearly half of LTCF residents, is more than 15 percent. In areas where there is a shortage of ICU beds and respirators, even the most carefully thought out ethical approaches to rationing these resources will place older patients at a lower priority. LTCFs must be prepared to manage patients who have had or have COVID-19 infection.

In an editorial published in the *Journal of the American Geriatrics Society*, Joseph G. Ouslander, M.D., an internationally renowned geriatrician, a professor of geriatric medicine, and a senior advisor to the dean at Florida Atlantic University's Schmidt College of Medicine, provides key updates and resources for front-line nursing home [staff](#) and clinicians.

"Until more rapid COVID testing is available to enable us to test all nursing home patients and staff, cases of COVID-19 infection are likely to continue to increase in the long-term care facility setting," said Ouslander. "To complicate matters, [long-term care facilities](#) will be asked to accept patients with possible or known COVID-19 from hospitals if the local or state government has not required the suspension of admissions during the presence of active COVID-19 cases."

Ouslander recommends developing an emergency plan that addresses patient/resident placement and staffing considerations before a cluster infection or outbreak occurs; employing intensive infection control practices; behaving as if all patients and staff are potentially infected with the virus; and continuing to carefully screen staff, recognizing that some may be asymptomatic carriers of the virus and pass the screening.

In addition to restricting visitors, communal dining and cancelling other

group activities, Ouslander advises providing therapy in the patient/resident room when feasible; if not, then safe social distancing should be used in therapy areas. Staff should wear plain surgical masks at all times and use N95 masks if available when performing high-risk procedures such as respiratory treatments. Full personal protective equipment (PPE) should be used when appropriate and available based on patient/resident symptoms and signs.

Staff who are quarantined based on symptoms and want to return to work must be tested to confirm they are not shedding the virus; returning health care workforce should be a priority for rapid testing. In addition, clinicians should monitor vital signs at least daily and more frequently if indicated; and even subtle changes in a patient's condition should be taken seriously and further evaluated when clinically indicated.

Because LTCF patients/residents cannot have visitors or participate in group activities, their isolation may have adverse effects on their [mental health](#). Morale of staff may also be affected. Experience in China suggests that half or more of [health care workers](#) treating patients with COVID-19 infection had symptoms of depression, anxiety, insomnia or distress. Ouslander suggests that LTCFs provide as much psychosocial care and support as feasible and use social media and video phone call options. Grief counseling for families and staff, as well as assistance with burial arrangements may be needed.

In the United States, of the approximately 1.2 million registered nurses employed outside of hospital settings, 24 percent are ages 55 to 64 years and 5 percent are age 65 years or older. Of the approximately 1.2 million physicians, an estimated 230,000 (20 percent) are ages 55 to 64 years and an estimated 106,000 (9 percent) are age 65 years or older. Many hospitals and health care systems are developing strategies to use these health professionals in productive ways that may not necessarily involve direct patient contact. The U.S. Centers for Medicare and Medicaid

Services has issued many waivers that could help bolster the health care workforce and the capacity of the nation's health care system to care for older COVID-19 patients.

"During this challenging and unprecedented pandemic of our lifetimes, we should be thankful for all of the front-line nursing home staff and clinicians who are risking their health and that of their families to care for the most vulnerable among us," said Ouslander. "Shortages of these professionals are anticipated, and the aging of our health care workforce makes many of us even more susceptible to acquiring COVID-19 and its complications. We all have a role to play, and we should support our geriatric health care workforce, our patients, and their families in whatever ways we can contribute to meet this daunting challenge."

**More information:** Joseph G. Ouslander, Coronavirus-19 in Geriatrics and Long-Term Care: An Update, *Journal of the American Geriatrics Society* (2020). [DOI: 10.1111/jgs.16464](https://doi.org/10.1111/jgs.16464)

Provided by Florida Atlantic University

Citation: Caring for those most vulnerable to COVID-19 requires vigilance (2020, April 16) retrieved 2 May 2024 from <https://medicalxpress.com/news/2020-04-vulnerable-covid-requires-vigilance.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--