

# App calculates risk of delaying cancer care during pandemic

May 26 2020

---



Credit: CC0 Public Domain

As the COVID-19 pandemic has overwhelmed healthcare systems across the country, doctors have postponed surgery and other treatments for thousands of patients with cancer. These delays may last for months, especially for those with early stage and less aggressive disease.

Unfortunately, there hasn't been a good way for doctors to compare the long-term risk to a patient from a months-long postponement of care to the additional risk posed by potential COVID-19 infection if they undergo surgery, chemotherapy and/or radiation.

So, a team of data scientists and cancer doctors from the University of Michigan Rogel Cancer Center and the U-M School of Public Health developed a free, web-based application to do just that.

The OncCOVID app draws on large, national cancer data sets to help assess the risk from of immediate treatment versus delayed treatment, depending on a patient's individual characteristics, as well as on COVID's impact on their local community.

"For many types of cancer, the data show delays in treatment lead to worse outcomes for patients," says the project's lead researcher, Holly Hartman, a doctoral student in biostatistics at U-M. "But each time a [cancer patient](#) goes to the hospital to receive care, they're also putting themselves at higher risk of contracting COVID-19.

So, it's essential to balance the need for treatment for this very serious disease and the extra risk that COVID-19 poses for cancer patients, whose immune systems are often compromised."

The researchers envision OncCOVID being used by doctors to help identify patients whose risk from COVID is outweighed by the benefits of immediate treatment.

"We also see the app providing additional reassurance to oncologists and their patients when the data show that delaying treatment will likely have little or no impact on a patient's long-term outcome," Hartman adds.

Meanwhile, OncCOVID could also be used by health care systems that are ramping services back up and need to prioritize a backlog of patients whose treatment was put on hold due to the pandemic, adds Daniel Spratt, M.D., an associate professor of radiation oncology at Michigan Medicine and one of the senior researchers on the project.

"Hospitals have basically been using a three-tiered system during COVID: treat, delay a little, or delay a lot," Spratt says. "Unfortunately, this tiered system is an extremely blunt instrument. Our goal was to create a resource that could be tailored both to the individual patient and to their [local community](#)."

The app allows doctors to enter more than 45 characteristics about a patient—including their age, location, cancer type and stage, treatment plan, underlying medical conditions, and the proposed length of a delay in care. It then calculates the patient's likely five-year survival following immediate treatment and delayed treatment.

Under the hood, the app draws on millions of records contained in the National Cancer Institute's Surveillance, Epidemiology, and End Results (SEER) registry and the National Cancer Database, combined with county-level COVID infection data from Johns Hopkins University.

In some cases, the personalized risk assessments run counter to the generic, three-tiered approach.

For example, Spratt says, the team's model shows that an otherwise healthy 45-year-old woman from Ann Arbor, Mich. with stage 1 breast cancer actually has a slightly higher risk of dying over the next five years

if treatment is delayed by more than three months than if she's treated immediately.

"Under the three-tiered model, care for this patient wouldn't be considered life-threatening and it's likely her care may be delayed for months under most of the published tier-based systems," he notes.

Conversely, under the team's data model, a 70-year-old woman from New York City—where COVID infections have been high—with stage 1 breast cancer and several underlying health conditions would be put at significantly more risk by undergoing immediate treatment than the risk posed by a three-month delay.

For the data savvy, advanced features allow all of the app's underlying statistical assumptions to be adjusted—such as the baseline mortality risk from COVID and the disease's infection rate.

In the future, the researchers plan to use the same data model to start looking at the effects that treatment delays due to the [coronavirus](#) pandemic may have on cancer mortality nationally, Hartman adds.

The OncCOVID app is not intended to provide medical advice to patients, the researchers caution. A number of factors may figure into a care provider's recommendation to delay or proceed with cancer treatment, including their local hospital's capacity to safely treat [cancer](#) patients during the pandemic.

Provided by University of Michigan

Citation: App calculates risk of delaying cancer care during pandemic (2020, May 26) retrieved 3 May 2024 from <https://medicalxpress.com/news/2020-05-app-cancer-pandemic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.