

Contact tracing can cut time to case isolation in SARS-CoV-2

May 1 2020



(HealthDay)—Contact tracing can reduce the time to case isolation of

severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), according to a study published online April 27 in *The Lancet Infectious Diseases*.

Qifang Bi, from Johns Hopkins Bloomberg School of Public Health in Baltimore, Maryland, and colleagues compared cases identified through symptomatic surveillance and [contact tracing](#) using data from 391 SARS-CoV-2 cases and 1,286 close contacts identified from Jan. 14 to Feb. 12, 2020. The time from [symptom onset](#) to confirmation, isolation, and admission to hospital was estimated.

The researchers found that 91 percent of 391 cases had mild or moderate clinical severity at initial assessment. Three cases had died and 225 had recovered as of Feb. 22, 2020 (median time to recovery, 21 days). On average, [cases](#) were isolated 4.6 days after symptom development; this was reduced by 1.9 days with contact tracing. The risk for infection was higher for household contacts and those traveling with a case (odds ratios, 6.27 and 7.06, respectively) compared with other close contacts. There was an 11.2 percent household secondary attack rate; children were as likely to be infected as adults (infection rate, 7.4 and 6.6 percent for children

Citation: Contact tracing can cut time to case isolation in SARS-CoV-2 (2020, May 1) retrieved 30 January 2023 from <https://medicalxpress.com/news/2020-05-contact-case-isolation-sars-cov-.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.