

## How and when to end the COVID-19 lockdown: An optimization approach

May 6 2020



Credit: CC0 Public Domain

Countries around the world are in a state of lockdown to help limit the spread of SARS-CoV-2. However, as the number of new daily confirmed cases begins to decrease, governments must decide how to release their populations from quarantine as efficiently as possible without overwhelming their health services.



In a preprint manuscript published this week, led by Professor Mike Bonsall from the Department of Zoology at Oxford University, new research applied an optimal control framework to an adapted Susceptible-Exposure-Infection-Recovery (SEIR) model framework to investigate the efficacy of two potential lockdown release strategies. In this instance the UK population was used as a test case.

Professor Bonsall said: "What we found was that to limit recurrent spread, ending quarantine for the entire population simultaneously is a high-risk strategy, and that a gradual re-integration approach would be more reliable.

"Furthermore, to increase the number of people that can be first released, lockdown should not be ended until the <u>number</u> of new daily confirmed cases reaches a sufficiently low threshold."

The team modeled a gradual release strategy by allowing different fractions of those in lockdown to re-enter the working non-quarantined population. Mathematical optimization methods, combined with an adapted SEIR model, determined how to maximize those working while preventing the health service from being overwhelmed.

The analyses showed the <u>optimal strategy</u> to be to release approximately as many as half the <u>population</u> two-to-four weeks from the end of an initial infection peak, then wait another three-to-four months to allow for a second peak before releasing everyone else.

Co-author Thom Rawson, from the Department of Zoology at Oxford University, said: "We also modeled an 'on-off' strategy of releasing everyone, but re-establishing lockdown if infections become too high. We found that the worst-case scenario of a gradual release is more manageable than the worst-case scenario of an on-off strategy, and caution against lockdown-release strategies based on a threshold-



dependent on-off mechanism. The potential for a large increase in infection numbers adds further evidence to the need for increased testing capacity in the wake of ending lockdown."

The two quantities most critical in determining the optimal solution are transmission rate and the recovery rate, where the latter is defined as the fraction of infected people in any given day that then become classed as recovered. The team suggest that the accurate identification of these values is of particular importance to the ongoing monitoring of the pandemic.

Co-author Professor Chris Huntingford, of the UK Centre for Ecology & Hydrology, who applied his mathematical expertise to the modeling, said: "As we understand more about COVID19, such numbers in the computer model can be adjusted and simulations repeated. Hence the optimum release strategy from lockdown can be updated if required, and as we learn more in the weeks ahead."

**More information:** Thomas Rawson et al. How and when to end the COVID-19 lockdown: an optimisation approach, (2020). <u>DOI:</u> 10.1101/2020.04.29.20084517

Provided by University of Oxford

Citation: How and when to end the COVID-19 lockdown: An optimization approach (2020, May 6) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2020-05-covid-lockdown-optimization-approach.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.