

How has the COVID-19 pandemic affected women's sexual behavior?

May 12 2020



Credit: CC0 Public Domain

A recent study from Turkey found that women's sexual desire and frequency of intercourse increased during the COVID-19 pandemic, but their quality of sexual life decreased. The findings are published in the

International Journal of Gynecology & Obstetrics.

In the study of 58 women, women participated in [sexual intercourse](#) on average 2.4 times per week during the pandemic, compared with 1.9 times in the 6–12 months prior to the pandemic. Before the pandemic 32.7% of participants desired to become pregnant, compared with 5.1% during the pandemic; however, use of contraception decreased during the pandemic.

Menstrual disorders were more common during the pandemic than before (27.6% versus 12.1%), and participants generated worse scores on a questionnaire based on sexual function during the pandemic compared with scores before the pandemic.

More information: Bahar Yuksel et al. Effect of the COVID-19 pandemic on female sexual behavior, *International Journal of Gynecology & Obstetrics* (2020). [DOI: 10.1002/ijgo.13193](https://doi.org/10.1002/ijgo.13193)

Provided by Wiley

Citation: How has the COVID-19 pandemic affected women's sexual behavior? (2020, May 12) retrieved 25 April 2024 from <https://medicalxpress.com/news/2020-05-covid-pandemic-affected-women-sexual.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.