

Diastolic dysfunction more dangerous than previously thought

May 11 2020, by Johannes Angerer



Credit: CC0 Public Domain

Sudden cardiac death is a common cause of death in patients with reduced systolic ejection function. As part of a long-term observational study, MedUni Vienna researchers from the Division of Cardiology have



now shown the importance of the diastolic filling function: the less the heart fills when there is diastolic dysfunction, the more likely patients are to die suddenly of cardiac arrhythmia.

People with ischemic (narrowing of the coronary arteries) or nonischemic myocardial disease have an inherently higher risk of <u>cardiac</u> <u>arrhythmia</u>. Where the heart's ejection function is significantly reduced (systolic heart failure), <u>current practice</u> is to also consider an implantable defibrillator (ICD). The ICD then emits an electrical pulse when there is rapid ventricular arrhythmia and/or ventricular fibrillation, until a normal heart rhythm is resumed.

However, every second heart failure patient suffers from the diastolic form (reduced filling function) of heart failure: the stiffened <u>left</u> <u>ventricle</u> cannot expand sufficiently and is filled against high resistance. This means that less blood can flow into the left ventricle but all of it is pumped into the circulatory system. It is even possible that the systolic ejection function (also called the ejection fraction) is maintained. Women and more <u>elderly people</u> are more likely to suffer from this form of heart failure.

"Hitherto, we have distinguished between these two forms when it comes to assessing the risk of <u>sudden cardiac death</u>," says Principal Investigator Thomas Pezawas from the Division of Cardiology, "Patients with nonsystolic cardiac insufficiency are assessed as having a lower risk of sudden cardiac death and are therefore rarely given a primary preventive ICD."

Diastolic dysfunction also dangerous

For the purposes of the study, 210 at-risk patients who did not previously have cardiac arrhythmia, or only had a mild form of it, were investigated over a period of up to 10 years. "The number of potentially fatal cases is



much higher than expected and the correlation with the level of <u>diastolic</u> <u>dysfunction</u> is striking," says Pezawas, describing the findings that have now been published in top journal *Circulation: Arrhythmia and Electrophysiology*. "Unfortunately, sudden cardiac death also affects patients with only mild cardiac insufficiency."

These results will also be important for other centers, since the data available in this field was previously very sparse, say the MedUni Vienna expert. The aim is to achieve an excellent level of protection against sudden cardiac death and to screen the right patients. The recommendation made by the study authors to include diastolic function in the risk assessment could bring about a paradigm shift in treatment.

More information: Thomas Pezawas et al. Importance of Diastolic Function for the Prediction of Arrhythmic Death, *Circulation: Arrhythmia and Electrophysiology* (2020). DOI: <u>10.1161/CIRCEP.119.007757</u>

Provided by Medical University of Vienna

Citation: Diastolic dysfunction more dangerous than previously thought (2020, May 11) retrieved 16 August 2024 from https://medicalxpress.com/news/2020-05-diastolic-dysfunction-dangerous-previouslythought.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.