


Fluoridation is not associated with increase in osteosarcoma

May 12 2020



Credit: CC0 Public Domain

The *Journal of Dental Research* published today the results of a study that demonstrated that community water fluoridation is not associated with increased risk of osteosarcoma.

More than sixty percent of the U.S. population have access to community [water](#) fluoridation, considered to be one of the most important public health policies of the twentieth century due to its reduction of tooth decay at the population level. Fluoride ingestion has been suggested as a possible risk factor for osteosarcoma based on a 1990 animal study. Six of the seven subsequent case-control studies in humans reported that fluoride in drinking water was not associated with osteosarcoma.

This study assessed whether living in a fluoridated community was a risk factor for osteosarcoma by performing a secondary data analysis using data collected from two separate, but linked studies. Patients for both Phase 1 and Phase 2 were selected from U.S. hospitals using a hospital-based matched case-control study design. For both phases, cases were patients diagnosed with osteosarcoma and controls were patients diagnosed with other bone tumors or non-neoplastic conditions.

In Phase 1, cases (N=209) and controls (N=440) were patients of record in the participating orthopedic departments from 1989-1993. In Phase 2, cases (N=108) and controls (N=296) were incident patients who were identified and treated by orthopedic physicians from 1994-2000. This analysis included all [patients](#) who met eligibility criteria on whom we had complete data on covariates, exposures, and outcome. Conditional logistic regression was used to estimate odds ratios (OR) and 95% confidence intervals (CI) for the association of community water fluoridation with osteosarcoma.

The adjusted OR, for osteosarcoma and ever-having lived in a fluoridated area for non-bottled water drinkers was 0.51(0.31—0.84), $p=0.008$. The same comparison adjusted OR for bottled water drinkers was 1.86 (0.54—6.41), $p=0.326$.

"These results indicate that residence in a fluoridated community is not

related to an increase in risk for osteosarcoma after adjusting for race, ethnicity, income, distance from the hospital, urban/rural living status, and drinking bottled water. This should not be surprising given that ingestion of fluoridated water is a common exposure and [osteosarcoma](#) remains a [rare disease](#)," said Chester Douglass; Harvard School of Dental Medicine, Department of Oral Health Policy and Epidemiology.

More information: F.M. Kim et al. A Case-Control Study of Fluoridation and Osteosarcoma, *Journal of Dental Research* (2020).
[DOI: 10.1177/0022034520919385](https://doi.org/10.1177/0022034520919385)

Provided by International & American Associations for Dental Research

Citation: Fluoridation is not associated with increase in osteosarcoma (2020, May 12) retrieved 7 May 2024 from <https://medicalxpress.com/news/2020-05-fluoridation-osteosarcoma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
