

Immunotherapy improves survival in patients with advanced bladder cancer

May 29 2020



Cancer cell during cell division. Credit: National Institutes of Health

An immunotherapy drug called 'avelumab' has been shown to significantly improve survival in patients with the most common type of bladder cancer, according to results from a phase III clinical trial led by

Professor Tom Powles from Queen Mary University of London and Barts Cancer Centre, UK.

This is the first time an [immune therapy](#) has resulted in a survival advantage in this setting in bladder cancer, and will potentially benefit thousands of patients each year.

The results will be presented in the plenary session at the American Society of Clinical Oncology, the world's largest annual cancer meeting (ASCO20). They found that avelumab led to a 31 per cent reduction in risk of death of bladder cancer and extended median survival in advanced bladder cancer by more than seven months.

Approximately 550,000 new cases of bladder cancer are diagnosed each year (10,200 of which are in the UK), making it the tenth most common cancer worldwide. This trial focused on the group of these patients whose cancer had spread beyond the [bladder](#) (advanced or stage 4 disease), which is difficult to treat.

Chemotherapy is the current initial standard of care in the treatment of these advanced cancers. After chemotherapy is finished, patients are checked regularly because the cancer tends to return quickly. When it returns it is difficult to treat and outcomes are poor.

The phase III global trial, named JAVELIN Bladder 100 and funded by Pfizer and Merck KGaA Darmstadt, Germany, evaluated the efficacy of the [immunotherapy drug](#) 'avelumab' in patients with locally advanced or metastatic urothelial carcinoma which they received after the initial chemotherapy.

A total of 700 patients from over 200 sites around the world were then assigned to two treatment groups after the completion of chemotherapy—one group receiving regular checking (standard care) on

its own, and the other receiving avelumab in addition to standard care.

Treatment with avelumab resulted in a 31 per cent reduction in risk of death and [median overall survival](#) of 21.4 months compared with 14.3 months in patients who did not receive the drug. Side effects were in line with expectations with immune therapy and 11 per cent of patients stopped avelumab due to treatment problems.

Study lead Thomas Powles, Professor of Genitourinary Oncology at Queen Mary University of London, and Director of Barts Cancer Centre, Barts Health NHS Trust, said: "This is the first time that an immune therapy clinical trial has shown a survival benefit for first-line therapy in metastatic [bladder cancer](#).

"We saw a meaningful reduction in the risk of death and a significant overall [survival benefit](#) with avelumab, which underscores the potential for this immunotherapy to be practice-changing for patients. This highlights the potential benefits of a maintenance approach with avelumab in patients to prolong their lives following chemotherapy."

Avelumab is a type of immunotherapy, known as a checkpoint inhibitor, which blocks a protein called PD-L1 on the surface of tumour cells. When PD-L1 is active it helps cancers hide from the immune system. By blocking PD-L1 the immune system finds it easier to identify and kill the [cancer](#).

More information: 'Maintenance avelumab + best supportive care (BSC) versus BSC alone after platinum-based first-line (1L) chemotherapy in advanced urothelial carcinoma (UC): JAVELIN Bladder 100 phase III interim analysis'. Thomas Powles, Se Hoon Park, Eric Voog, Claudia Caserta, B.P. Valderrama, Howard Gurney, Haralabos Kalofonos, Sinisa Radulovic, Wim Demey, Anders Ullén, Yohann Loriot, Srikala S. Sridhar, Norihiko Tsuchiya, Evgeny

Kopyltsov, Cora N. Sternberg, Joaquim Bellmunt, Jeanny B. Aragon-Ching, Daniel Peter Petrylak. American Society of Clinical Oncology ASCO20 conference Plenary Session.

meetinglibrary.asco.org/record/186872/abstract

Provided by Queen Mary, University of London

Citation: Immunotherapy improves survival in patients with advanced bladder cancer (2020, May 29) retrieved 2 May 2024 from <https://medicalxpress.com/news/2020-05-immunotherapy-survival-patients-advanced-bladder.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--