

Isolating the ill and prioritising remote work are key strategies in combating the coronavirus

May 27 2020



Credit: CC0 Public Domain

In March of this year, Aalto University, the Finnish Meteorological Institute, VTT Technical Research Centre of Finland and the University

of Helsinki launched a joint project aimed at investigating airborne transmission and spread of coronavirus in indoor spaces.

When a person speaks, cough or sneezes, droplets are generated from their [respiratory tract](#), and these can carry pathogens such as coronaviruses.

Researchers have now published the first, preprint version of the paper, which has been submitted for peer-review and published at Arxiv.org. The paper details how they have modelled the airborne transport of different-sized droplets. These are emitted through coughing, so the study evaluated the quantities of particles that someone could come into contact with upon entering a supermarket or any other indoor public space.

Assistant professor at Aalto University, and project coordinator, Ville Vuorinen, says that both previous related research, and a number of well-known [infection](#) spikes, indicate a substantial risk of coronavirus through inhalation of aerosol particles, as well as direct droplet transmission and transmission from surfaces. The 3-D flow simulations and analyses carried out in the project also support these ideas.

"Our simulations show that virtually all droplets of less than 50 micrometres—and thus the majority of those produced when coughing—dry up as particles before they reach the floor and then linger around as carried by the indoor airflows. We consider it possible that these particles contain enough virus pathogens to cause infection," says Vuorinen.

Docent Antti Hellsten, from the Finnish Meteorological Institute, says "We modelled a typical indoor space, separated with shelves and room dividers, with high-resolution 3-D flow simulations. In the modeling, the coughing of a person carrying the coronavirus raised the particle content

in the immediate environment to such a high level that risk for exposure was still considered to be significant about four metres away. The risk remained significant for up to several minutes afterwards."

Although the risk of infection during a single shopping trip is relatively low, Vuorinen emphasises that at the probability of infection accumulates with regular exposure throughout several weeks.

"Tens of millions of grocery trips are carried out in Finland each month, and the likelihood of exposure in a grocery store could be as high as 1/1000 during particular stages of the pandemic. This would mean that as many as 10 000 Finns are exposed each month. If infectious people with symptoms stay at home and people reduce their excursions to public places, the risk decreases significantly," Vuorinen explains.

Even talking can be enough to transmit the virus

For the paper, the researchers examined well-known infection spikes in which a large number of people are believed to have been infected by a single pre-symptomatic infectious person. Based on other similar cases and previous studies, the researchers estimated that participants could have inhaled 50-500 aerosol particles which could have been enough for significant exposure.

"In the modeling, we examined the effect of the length of time spent in the space, the total number of people and the number of those coughing. One person per square metre, as is common in bars, just simply talking, can create a particle cloud sufficient for infecting others nearby. A symptomatic person working in an open office can create concentrations in the surrounding space ranging, on average, from 10 to 500 aerosol particles per cubic metre. If 100 aerosol particles per cubic metre are taken as the risk limit for exposure, people may therefore have exposure in times ranging from a few minutes to as long as several hours,"

Vuorinen adds.

According to the researchers, it is likely that infection spikes always involve direct contact infection, aerosol infection and shared surface infections. The longer you are there, and the closer the proximity to others, the greater the risk of infection.

Isolation measures have gradually started to be relaxed in Finland and many other countries, but researchers still recommend doing work remotely as much as possible. That being said, the safety of workplaces could be improved in many ways.

"The most important thing is that nobody comes to work sick. There should be the appropriate ventilation, sufficient space between workers and with only the minimum amount of personnel necessary. This reduces both people's proximity to each other and possible exposure time," Vuorinen explains.

Supercomputer used for modelling tasks

The project involves around 30 researchers whose specialisations include flow dynamics, aerosol physics, social networks, ventilation, biomedical engineering, [health sciences](#) and medicine.

The airborne movement and preservation of droplets leaving the respiratory tract were calculated using the supercomputer owned by CSC—IT Center for Science Ltd, and various analyses and 3-D visualisations of the results were carried out.

"The physics of the phenomena now modelled are already reasonably well known, but this new research sheds new light on them, particularly from the perspective of the coronavirus and public spaces," says Ville Vuorinen.

"Our understanding of the airborne transmission of SARS-CoV-2 virus has changed significantly in recent months," says Assistant Professor Tarja Sironen from University of Helsinki.

More information: Modelling aerosol transport and virus exposure with numerical simulations in relation to SARS-CoV-2 transmission by inhalation indoors: arxiv.org/abs/2005.12612

Provided by Aalto University

Citation: Isolating the ill and prioritising remote work are key strategies in combating the coronavirus (2020, May 27) retrieved 30 April 2024 from <https://medicalxpress.com/news/2020-05-isolating-ill-prioritising-remote-key.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--