


Pediatric coronavirus disease (COVID-19) X-ray, CT in review of new lung disorders

May 9 2020



Reverse transcription-polymerase chain reaction testing confirmed diagnosis of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). A, Frontal chest radiograph obtained at initial presentation shows bilateral lower lung zone-predominant consolidations and, to lesser extent, ground-glass opacities. B, Frontal chest radiograph obtained 2 days after hospital admission shows interval increase in consolidation in bilateral lower lung zones. C, Frontal chest radiograph obtained 6 days after hospital admission and treatment shows interval improvement in consolidations in bilateral lower lung zones. Credit: *American Journal of Roentgenology* (AJR)


Although the clinical symptoms of new pediatric lung disorders such as severe acute respiratory syndrome (SARS), swine-origin influenza A (H1N1), Middle East respiratory syndrome (MERS), e-cigarette or vaping product use-associated lung injury (EVALI), and coronavirus

disease (COVID-19) pneumonia may be nonspecific, some characteristic imaging findings "have emerged or are currently emerging," according to an open-access article in the *American Journal of Roentgenology* (AJR).

"Although there are some overlapping imaging features of these disorders," wrote first author Alexandra M. Foust of Boston Children's Hospital and Harvard Medical School, "careful evaluation of the distribution, lung zone preference, and symmetry of the abnormalities with an eye for a few unique differentiating imaging features, such as the halo sign seen in COVID-19 and subpleural sparing and the atoll sign seen in EVALI, can allow the radiologist to offer a narrower differential diagnosis in [pediatric patients](#), leading to optimal patient care."

At most institutions, whereas the first imaging study performed in patients with clinically suspected COVID-19 is [chest](#) radiography, Foust and colleagues' review of the clinical literature found that studies on chest radiography findings in patients with COVID-19 were relatively scarce.

Regarding the limited studies of pediatric patients with COVID-19, Foust et al. noted chest radiography "may show normal findings; patchy bilateral ground-glass opacity (GGO), consolidation, or both; peripheral and lower lung zone predominance."



Reverse transcription-polymerase chain reaction testing confirmed diagnosis of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). A-D, Axial (A and B), coronal (C), and sagittal (D) lung window setting CT images show posterior subpleural ground-glass opacity with small component of consolidation within left lower lobe. Credit: *American Journal of Roentgenology* (AJR)

Similarly, while the literature describing chest CT findings in patients with COVID-19 are more robust than those describing chest radiography findings, only a few articles have reported CT findings of COVID-19 in children.

A study of 20 pediatric patients with COVID-19 reported that the most frequently observed abnormalities on CT were subpleural lesions (100% of patients), unilateral (30%) or bilateral (50%) pulmonary lesions, GGO (60%), and consolidation with a rim of GGO surrounding it, also known as the halo sign (50%).

The authors of this AJR article also pointed to a smaller study of five pediatric patients with COVID-19, where investigators reported modest patchy GGO, one with peripheral subpleural involvement, in three patients that resolved on follow-up CT examination.

More information: Alexandra M. Foust et al, Pediatric SARS, H1N1, MERS, EVALI, and Now Coronavirus Disease (COVID-19) Pneumonia: What Radiologists Need to Know, *American Journal of Roentgenology* (2020). [DOI: 10.2214/AJR.20.23267](https://doi.org/10.2214/AJR.20.23267)

Provided by American Roentgen Ray Society

Citation: Pediatric coronavirus disease (COVID-19) X-ray, CT in review of new lung disorders (2020, May 9) retrieved 26 April 2024 from <https://medicalxpress.com/news/2020-05-pediatric-coronavirus-disease-covid-x-ray.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.