

Antioxidants in brain linked to improved psychosis treatment

June 9 2020



Researchers studied antioxidant levels in an area of the brain called cingulate cortex which is well connected to a network of regions that play a major role in generating symptoms of psychosis. Credit: University of Western Ontario

Once patients with psychosis start treatment, some get better in weeks while it can take months for others.

Seeking to understand and influence this disparity, Schulich School of Medicine & Dentistry professor Dr. Lena Palaniyappan and his team are exploring specific chemicals in the brain with the aim of speeding up the time it takes a patient to respond to medication.



Past research has shown that patients who experience their first episode of psychosis and respond early to treatment have better overall outcomes. This includes improved symptoms and daily functioning as well as spending less time in hospital.

The research team from Schulich and Lawson Health Research Institute looked specifically at <u>antioxidant levels</u> in the brain and found that these chemicals which rid the body of normal metabolic biproducts, called free radicals, may improve outcomes of early intervention in psychosis. They looked specifically at an area of the brain called <u>cingulate cortex</u> which is well connected to a network of regions that play a major role in generating symptoms of psychosis.

The findings showed that patients with higher levels of an antioxidant called glutathione responded more quickly to medication for psychosis and had improved outcomes. They estimated that a 10 percent increase in antioxidants could lead to a reduction in length of hospital stay by at least seven days.

"This study demonstrates that if we can find a way to boost the amount of antioxidants in the brain, we might be able to help patients transition out of hospital more quickly, reduce their suffering more quickly and help them return earlier to their work and studies," said Palaniyappan, the Tanna Schulich Endowed Chair in Neuroscience and Mental Health at Western.

Antioxidant levels in the brain vary naturally from person to person and those variations can be attributed to lifestyle choices like exercise and diet. There are also ways to pharmaceutically boost these levels. A supplement called N-Acetyl Cysteine (NAC) as well as broccoli extracts can increase the <u>brain</u>'s antioxidants if given over a long period of time.

Palaniyappan and his team in collaboration with Dr. Jean Theberge at



Lawson Imaging used high-field MRI imaging to examine antioxidant levels in the brains of 37 new patients to the Prevention and Early Intervention for Psychoses Program (PEPP) at London Health Sciences Centre. Antioxidant levels were studied prior to beginning treatment for psychosis and followed up for six months post-treatment.

The study, "Early treatment response in first episode psychosis: a 7-T <u>magnetic resonance</u> spectroscopic study of glutathione and glutamate," was published in *Nature Molecular Psychiatry*.

More information: Kara Dempster et al. Early treatment response in first episode psychosis: a 7-T magnetic resonance spectroscopic study of glutathione and glutamate, *Molecular Psychiatry* (2020). DOI: 10.1038/s41380-020-0704-x

Provided by University of Western Ontario

Citation: Antioxidants in brain linked to improved psychosis treatment (2020, June 9) retrieved 27 June 2024 from https://medicalxpress.com/news/2020-06-antioxidants-brain-linked-psychosis-treatment.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.