

Campylobacter infection major cause of child stunting

June 5 2020, by Claudia Caruana



An infant undergoes medical screening. Stunting is a common issue among children in Bangladesh. Credit: [Maggie Moore/USAID \(CC BY-NC 2.0\)](#)

Gastrointestinal infection with *Campylobacter* plays a major role in the stunting of children in urban Bangladesh, say the authors of a new study.

The bacteria *Campylobacter*, which people get from contaminated food and drinks, can lead to severe diarrhoea. Infections of *Campylobacter* in

children under the age of two years are especially frequent in developing countries. According to a WHO factsheet, *Campylobacter* diarrhoea's duration and possible complications makes it highly important from a socio-economic perspective.

Stunting may result in neurodevelopmental delays and metabolic disorders later in life. It is a major global health challenge particularly in South Asia where 35 percent of all children are stunted. Worldwide, approximately 144 million children are stunted, according to the UN Children's Fund.

"Enteric infection, or [gastrointestinal infection](#), which includes *Campylobacter* infection, is associated with stunting through its effect on appetite, metabolic demands, and gastrointestinal absorption and function," says J. Johanna Sanchez, researcher at the Children's Health Research Centre, University of Queensland, Brisbane, Australia, and an author of the study published May in *PLOS Neglected Tropical Diseases*.

The study examined the linear growth of 265 children from 0 to 24 months of age, factoring in *Campylobacter* infection and household conditions that may have affected their growth. In the population studied, 70 percent of children under two years of age had *Campylobacter* infection.

"This is very important given the negative effect of enteric infection. Also, given the high global prevalence, this is a very important public health challenge to address. Although [environmental factors](#) resulting in *Campylobacter* infection may differ in different regions, studies say that the stunting effect is similar," says Sanchez.

The researchers also found treating drinking water and increased antibiotic use reduced *Campylobacter* infection and therefore indirectly had a positive effect on growth. "But rather than suggest that [children](#)

take antibiotics as a way to improve growth," she cautions, "we see this as another example of infection, particularly *Campylobacter* [infection](#), having an effect on linear growth."

Tim Julian, group leader of Pathogens and Human Health at the Swiss Federal Institute of Aquatic Science and Technology, tells SciDev.Net that Sanchez and her co-researchers "affirm a relationship between *Campylobacter* infections and child growth and identify a narrow presumptive age range (12—18 months) most impacted by *Campylobacter* spp. infections. They also identify key relationships between infections and potential beneficial interventions, including antibiotic use and drinking [water treatment](#)".

An integrated assessment of the findings highlights the potential for targeted interventions to provide the largest gains in child health, specifically, safe drinking water treatment within a six-month window of highest susceptibility, he says.

Julian says that it is equally important that the researchers had repeated their study method on other important enteric pathogens, *E. coli* and *Giardia lamblia*, to find "no significant associations between asymptomatic carriage of these enteric pathogens and linear growth". This, he said, was additional evidence that reducing *Campylobacter* transmission was key to improving child health and well-being.

Julian stresses that the additional evidence of the importance of *Campylobacter* relative to other enteric pathogens on linear child growth helps focus efforts on reducing transmission so that child health and well-being can be improved.

Stunting is a complex problem related to conditions of poverty, particularly in low- and middle-income countries, concludes Sanchez. "This includes inadequate nutrition and poor environmental conditions

that may result in infectious disease, which negatively affects linear growth."

More information: J. Johanna Sanchez et al. Campylobacter infection and household factors are associated with childhood growth in urban Bangladesh: An analysis of the MAL-ED study, *PLOS Neglected Tropical Diseases* (2020). [DOI: 10.1371/journal.pntd.0008328](https://doi.org/10.1371/journal.pntd.0008328)

Provided by SciDev.Net

Citation: Campylobacter infection major cause of child stunting (2020, June 5) retrieved 5 May 2024 from <https://medicalxpress.com/news/2020-06-campylobacter-infection-major-child-stunting.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--