

Depression and anxiety rise among new moms amidst the COVID-19 pandemic

June 19 2020



Credit: CC0 Public Domain

Pregnant and postpartum women are already at a high risk of depression and anxiety—one in seven women struggle with symptoms in the perinatal period. But the coronavirus pandemic is exacerbating those

struggles according to a recent study published in *Frontiers in Global Women's Health*, which found that the likelihood of maternal depression and anxiety has substantially increased during the health crisis.

"The social and physical isolation measures that are critically needed to reduce the spread of the virus are taking a toll on the physical and [mental health](#) of many of us," says Dr. Margie Davenport of the University of Alberta, Canada, who co-authored the study.

For new moms, those stresses come with side effects. "We know that experiencing depression and anxiety during pregnancy and the postpartum period can have detrimental effects on the mental and [physical health](#) of both mother and baby that can persist for years." Such effects can include premature delivery, reduced mother-infant bonding, and developmental delays in infants.

The study surveyed 900 women—520 of whom were pregnant and 380 of whom had given birth in the past year—and asked about their depression and anxiety symptoms before and during the pandemic. Before the pandemic began, 29% of those women experienced moderate to high anxiety symptoms, and 15% experienced depressive symptoms. During the pandemic, those numbers increased—72% experienced anxiety and 41% experienced depression.

Because lockdown measures have affected [daily routines](#) and access to gyms, researchers also asked women whether their exercise habits had changed. Of the women surveyed, 64% reduced their [physical activity](#) since the pandemic began, while 15% increased and 21% experienced no change. Exercise is a known way to ease depression symptoms, so limited physical activity may result in an uptick in depressive symptoms. Indeed, the study found that women who engaged in at least 150 minutes of moderate physical activity a week had significantly lower symptoms of depression and anxiety.

The findings are somewhat limited given the fact that researchers could not survey women before the pandemic began (since they could not know a pandemic would occur). The women surveyed could only offer their pre-pandemic symptoms in hindsight. Also, while the researchers asked women about their symptoms using validated measures, only mental health care professionals can validly diagnose an individual with [depression](#) or anxiety.

The study was specifically interested in the impact of COVID-19 on new moms, but Davenport says maternal mental health is a critical issue no matter the time. "Even when we are not in a global pandemic, many pregnant and [postpartum women](#) frequently feel isolated whether due to being hospitalized, not having family or friends around or other reasons," she says.

"It is critical to increase awareness of the impact of social (and physical) isolation on the mental health of pregnant and postpartum [women](#)," Davenport continues. Increased awareness makes diagnosis and treatment—the ultimate goal—more likely.

More information: Margie H. Davenport et al, Moms Are Not OK: COVID-19 and Maternal Mental Health, *Frontiers in Global Women's Health* (2020). [DOI: 10.3389/fgwh.2020.00001](https://doi.org/10.3389/fgwh.2020.00001)

Provided by Frontiers

Citation: Depression and anxiety rise among new moms amidst the COVID-19 pandemic (2020, June 19) retrieved 16 June 2024 from <https://medicalxpress.com/news/2020-06-depression-anxiety-moms-covid-pandemic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.