


New imaging method tracks brain's elusive networks

June 10 2020



Brain signals are measured with high-density EEG. The signal source is simulated and estimated using artificial intelligence. Results are validated with clinical findings. Credit: Carnegie Mellon University College of Engineering

Understanding the source and network of signals as the brain functions is a central goal of brain research. Now, Carnegie Mellon engineers have created a system for high-density EEG imaging of the origin and path of normal and abnormal brain signals.

Bin He, head of the Carnegie Mellon University's Department of Biomedical Engineering, and colleagues are working on a core initiative of the National Institutes of Health, called Brain Research through Advancing Innovative Neurotechnologies (BRAIN).

The group's work with high-density EEG, reported in the journal *Nature Communications*, takes a major step toward realizing one of the central goals of the BRAIN initiative: "produce a revolutionary new dynamic picture of the brain that, for the first time, shows how [individual cells](#) and complex [neural circuits](#) interact in both time and space." In other words, develop a way to watch how the brain does what it does.

"Electroencephalography (EEG), has been used for decades to track brain signals," explained Shumin Wang, Ph.D., director of the NIBIB program in Bio-Electromagnetic Technologies. "Dr. He and his team developed a more powerful, high-density version of EEG that can track brain signals across much larger areas of the brain than previously possible. Artificial intelligence is then used to identify where these stronger signals originate and travel through the brain, with great accuracy."

The research team tested the high-density EEG system in patients with epilepsy. Individuals experience seizures caused by errant pulses of brain activity, known as the epileptogenic zone. Many patients' seizures can be controlled with medications. For those who are medication-resistant, surgically removing the epileptogenic zone is a clinical option. The ability of high-density EEG to accurately identify the epileptic source would allow for much-improved brain surgeries to efficiently remove only the problem area and spare surrounding brain tissue.

The team tested their method, termed FAST-IRES for spatio-temporal iteratively reweighted edge sparsity, on 36 epilepsy patients undergoing pre-operative testing. FAST-IRES non-invasive [brain](#) recordings were obtained over several days using a special high-density EEG array. The fact that FAST-IRES is non-invasive is extremely important, given that the standard pre-operative testing requires invasive EEG, which is necessary to pinpoint the epileptogenic region but comes with increased risks of infection, complications, and costs.

Across the 36 patients, the FAST-IRES method analyzed more than 1000 EEG spikes and 86 seizures occurred. The high-density recordings were then analyzed using the [artificial intelligence](#) piece of the FAST-IRES method and compared with the standard invasive EEG pre-operative testing and the surgical findings.

The FAST-IRES method was extremely accurate in identifying the position and extent of the epileptogenic zone in the patients, which was confirmed with the surgical data. "Our results clearly demonstrated that FAST-IRES could identify the epileptogenic zone with high precision, using non-invasive high-density EEG scalp recordings," said He. Collaborators on the study at the Mayo clinic are considering implementing the system in the future.

The team believes the FAST-IRES technique could be used for diagnosis and treatment of Alzheimer's, Parkinson's, stroke and even depression. "It is extremely satisfying to see our work make a significant impact in fulfilling one of the central goals of the BRAIN initiative," said He. "In addition to treating patients, we look forward to our work helping researchers make significant advances in our understanding of human neuroscience."

More information: Abbas Sohrabpour et al. Noninvasive electromagnetic source imaging of spatiotemporally distributed epileptogenic brain sources, *Nature Communications* (2020). [DOI: 10.1038/s41467-020-15781-0](https://doi.org/10.1038/s41467-020-15781-0)

Provided by National Institutes of Health

Citation: New imaging method tracks brain's elusive networks (2020, June 10) retrieved 23 July 2024 from <https://medicalxpress.com/news/2020-06-imaging-method-tracks-brain-elusive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.