

New study shows how ketamine combats depression

June 1 2020



Credit: CC0 Public Domain

In low doses, the anaesthetic drug ketamine has been shown to have a rapid effect on difficult-to-treat depression. Researchers at Karolinska Institutet report that they have identified a key target for the drug:



specific serotonin receptors in the brain. Their findings, which are published in *Translational Psychiatry*, offer the hope of new, more effective antidepressants.

Depression is the most common psychiatric diagnosis in Sweden, affecting one in 10 men and one in five women at some point during their lives. Between 15 and 30% of patients are not helped by the first two attempts at therapy, in which case, depression is designated difficult to treat. Studies have shown that low doses of the anaesthetic drug ketamine act rapidly on certain sufferers, but exactly how it works is unknown. A <u>nasal spray</u> containing ketamine has recently been approved in the U.S. and EU for patients with <u>treatment-resistant depression</u>.

Researchers at Karolinska Institutet in Sweden imaged the brains of study participants using a <u>positron emission tomography</u> (PET) camera in connection with ketamine treatment.

"In this, the largest PET study of its kind in the world, we wanted to look at not only the magnitude of the effect, but also whether ketamine acts via serotonin 1B receptors," says the study's first author Mikael Tiger, a researcher at the Department of Clinical Neuroscience, Karolinska Institutet. "We and another research team were previously able to show a low density of serotonin 1B receptors in the brains of people with depression."

In the first phase of the study, 30 people with difficult-to-treat depression were randomly assigned to either a ketamine-infusion group (20 individuals) or a placebo (saline) group. It was a randomized doubleblind study, so neither patient nor doctor initially knew who received the active substance. The participants' brains were imaged with a PET camera before the infusion and 24-72 hours afterwards.

In the next phase, those who so wished (29 individuals) received



ketamine twice a week for two weeks. The result was that over 70% of those treated with ketamine responded to the drug according to a rating scale for depression.

Serotonin plays a key role in depression and low levels are thought to be linked to more serious disease. There are 14 different kinds of receptor for this neurotransmitter on the surface of neurons. For their PET imaging, the researchers used a radioactive marker that binds specifically to serotonin 1B receptors. They found that the ketamine operated via these receptors in a formerly unknown mechanism of action. Binding to this receptor reduces the release of serotonin, but increases that of another neurotransmitter called dopamine. Dopamine is part of the brain's reward system and helps people to experience positive feelings about life, something that is often lacking in <u>depression</u>.

"We show for the first time that ketamine treatment increases the number of <u>serotonin</u> 1B <u>receptors</u>," says the study's last author Johan Lundberg, research group leader at the Department of Clinical Neuroscience, Karolinska Institutet. "Ketamine has the advantage of being very rapid-acting, but at the same time, it is a narcotic-classed drug that can lead to addiction. So it'll be interesting to examine in future studies if this receptor can be a target for new, effective drugs that don't have the adverse effects of <u>ketamine</u>."

More information: Mikael Tiger et al, A randomized placebocontrolled PET study of ketamine's effect on serotonin1B receptor binding in patients with SSRI-resistant depression, *Translational Psychiatry* (2020). DOI: 10.1038/s41398-020-0844-4

Provided by Karolinska Institutet



Citation: New study shows how ketamine combats depression (2020, June 1) retrieved 12 May 2024 from <u>https://medicalxpress.com/news/2020-06-ketamine-combats-depression.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.