

# Use loss of taste and smell as key screening tool for COVID-19, researchers urge

June 4 2020

---



Credit: CC0 Public Domain

King's College London researchers have called for the immediate use of additional COVID-19 symptoms to detect new cases, reduce infections and save lives.

In a letter published in *The Lancet*, the team discussed how loss of [taste](#)

and [smell](#)—anosmia—should form part of screening measures for the virus.

They said: "As countries slowly emerge from lockdown measures, it is imperative to correctly contact-trace infected individuals. We believe that having added [loss of smell](#) and taste to the list of COVID-19 symptoms is of great value as it will help tracing almost 16% of cases that otherwise would have been missed."

The team, led by Professor Tim Spector, previously reported that loss of smell and taste is a key predictor of COVID-19 in addition to the most established symptoms of a high temperature and a new, continuous cough. The relative importance of the extra symptom was however disputed by sections of the UK government when it announced it was including the extra symptom. This additional analysis of the COVID Symptom Study app data and its 3.7 million users sought to quantify the clinical value of recording loss of smell in the population.

From 76,260 people with symptoms who tested positive for COVID-19 up to 19 May, 28.5% never reported any fever or cough and 16% reported loss of smell but not fever or cough. The prevalence of loss of smell and taste was three-fold higher in individuals testing positive (65%) than in those testing negative (22%), the strongest single predictor of being infected, suggesting that people with loss of smell and taste should self-isolate for at least seven days or until they can be tested.

Professor Tim Spector from King's College London said: "We believe that loss of smell and taste is a very common COVID-19 symptom and in fact, occurs more often than fever and lasts longer (5 days on average compared to only 2 for fever). Infections could be reduced, and lives saved now that this non-flu-like [symptom](#) is widely recognised, and actions are taken."

The researchers suggest that policymakers should consider these findings and their implications for mass screening as part of other public health measures in key areas such as schools, hospitals, airports and [care homes](#).

Professor Spector said: "Our data suggests that low-cost so-called 'smell the difference' [screening tests](#), that are already being used in some workplaces to screen people as they enter buildings, would capture a larger number of positive cases than temperature sensors do. We therefore feel that it should form part of a wider public health approach to reducing the infection rate."

Provided by King's College London

Citation: Use loss of taste and smell as key screening tool for COVID-19, researchers urge (2020, June 4) retrieved 5 May 2024 from <https://medicalxpress.com/news/2020-06-loss-key-screening-tool-covid-.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--