

Mozart may reduce seizure frequency in people with epilepsy

June 10 2020



Credit: Pixabay/CC0 Public Domain

A new clinical research study by Dr. Marjan Rafiee and Dr. Taufik Valiante of the Krembil Brain Institute at Toronto Western Hospital, part of University Health Network, has found that a Mozart composition may

reduce seizure frequency in patients with epilepsy.

The results of the research study, "The Rhyme and Rhythm of Music in Epilepsy," was recently published in the international journal *Epilepsia Open*. It looks at the effects of the Mozart melody, "Sonata for Two Pianos in D Major, K. 448" on reducing seizures, as compared to another auditory stimulus—a scrambled version of the original Mozart composition, with similar mathematical features, but shuffled randomly and lacking any rhythmicity.

"In the past 15 to 20 years, we have learned a lot about how listening to one of Mozart's compositions in individuals with [epilepsy](#) appears to demonstrate a reduction in [seizure frequency](#)," says Dr. Marjan Rafiee, lead author on the study. "But, one of the questions that still needed to be answered was whether individuals would show a similar reduction in [seizure](#) frequency by listening to another [auditory stimulus](#)—a control piece—as compared to Mozart."

The researchers recruited 13 patients to participate in the novel, year-long study. After three months of a baseline period, half of the patients listened to Mozart's Sonata once daily for three months, then switched to the scrambled version for three months. The others started the intervention by listening to the scrambled version for three months, then switched to daily listening of Mozart.

Patients kept "seizure diaries" to document their seizure frequency during the intervention. Their medications were kept unchanged during the course of the study.

"Our results showed daily listening to the first movement of Mozart K.448 was associated with reducing seizure frequency in adult individuals with epilepsy," says Dr. Rafiee. "This suggests that daily Mozart listening may be considered as a supplemental therapeutic option

to reduce seizures in individuals with epilepsy."

Epilepsy is the most common serious neurological disorder in the world, affecting approximately 300,000 Canadians and 50 million people worldwide.

Many experience debilitating seizures. The treatment is often one or more anti-seizure medications. But for 30 per cent of patients, the medications are not effective in controlling their seizures.

"As a surgeon, I have the pleasure of seeing individuals benefit from surgery, however I also know well those individuals for whom surgery is not an option, or those who have not benefitted from surgery, so, we are always looking for ways to improve symptom control, and improve quality of life for those with epilepsy," says Dr. Taufik Valiante, senior author of the study and the Director of the Surgical Epilepsy Program at Krembil Brain Institute at UHN and co-Director of CRANIA.

"Like all research, ours raises a lot of questions that we are excited to continue to answer with further research and support from the epilepsy community."

While these results are promising, the next step is to conduct larger studies with more patients, over a longer period of time.

More information: Marjan Rafiee et al, Daily listening to Mozart reduces seizures in individuals with epilepsy: A randomized control study, *Epilepsia Open* (2020). [DOI: 10.1002/epi4.12400](https://doi.org/10.1002/epi4.12400)

Provided by University Health Network

Citation: Mozart may reduce seizure frequency in people with epilepsy (2020, June 10) retrieved 8 May 2024 from

<https://medicalxpress.com/news/2020-06-mozart-seizure-frequency-people-epilepsy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.