

Newborn respiratory distress up with maternal antidepressant use

June 8 2020



(HealthDay)—Risk for newborn respiratory distress increases in a dose-

response manner in association with maternal use of prenatal antidepressants, according to a study published online June 8 in *Pediatrics*.

Gretchen Bandoli, Ph.D., from the University of California San Diego in La Jolla, and colleagues identified 226,932 singleton deliveries from OptumLabs Data Warehouse to estimate the risk for neonatal outcomes from patterns of prenatal antidepressant use.

A total of 15,041 pregnancies (6.6 percent) were exposed to an antidepressant. Use patterns were classified as low use with first-trimester reduction, low sustained use, moderate use with first-trimester reduction, moderate sustained use, and high sustained use. The researchers found that the risk for major cardiac malformations was increased with moderate sustained use; when compared with depression or anxiety reference groups, the results included the null. Moderate and high sustained trajectories were associated with an [increased risk](#) for [preterm birth](#) (adjusted risk ratios, 1.31 and 1.78, respectively). The risk for neonatal respiratory distress was increased with all four trajectories in a dose-response manner (adjusted risk ratios, 1.36 to 2.23) compared with the lowest trajectory as the reference group.

"Our findings support the continued use of the methodology to further delineate risk by different patterns of antidepressant use," the authors write. "This approach can help clinicians counsel [pregnant women](#) on the use of antidepressants during gestation."

More information: [Abstract/Full Text](#)
[Editorial](#)

Copyright © 2020 [HealthDay](#). All rights reserved.

Citation: Newborn respiratory distress up with maternal antidepressant use (2020, June 8)
retrieved 11 May 2024 from <https://medicalxpress.com/news/2020-06-newborn-respiratory-distress-maternal-antidepressant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.