


New study looks at post-COVID-19 emerging disease in children

June 25 2020



Thoracic findings in a 15-year-old girl with Multisystem Inflammatory

Syndrome in Children (MIS-C). (a) Chest radiograph on admission shows mild perihilar bronchial wall cuffing. (b) Chest radiograph on the third day of admission demonstrates extensive airspace opacification with a mid and lower zone predominance. (c, d) Contrast-enhanced axial CT chest of the thorax at day 3 shows areas of ground-glass opacification (GGO) and dense airspace consolidation with air bronchograms. (c) This conformed to a mosaic pattern with a bronchocentric distribution to the GGO (white arrow, d) involving both central and peripheral lung parenchyma with pleural effusions (black small arrow, d). Credit: Radiological Society of North America

In recent weeks, a multisystem hyperinflammatory condition has emerged in children in association with prior exposure or infection to SARS-CoV-2. A new case series published in the journal *Radiology* examines the spectrum of imaging findings in children with the post-COVID-19 inflammatory condition known in the U.S. as Multisystem Inflammatory Syndrome in Children (MIS-C).


The array of findings includes airway inflammation and rapid development of pulmonary edema, coronary artery aneurysms, and extensive intra-abdominal inflammatory changes.

In April 2020, Evelina London Children's Hospital in London, U.K., experienced a surge of [children](#) with a multi-system hyperinflammatory syndrome. The children had a variety of symptoms, including fever, headaches, abdominal pain, rash and conjunctivitis. Clinical features and lab findings shared some similarities to those of Kawasaki disease—which causes inflammation in the walls of blood vessels—Kawasaki-disease shock syndrome or toxic-shock syndrome, although atypical and more severe.

"Our hospital saw an unprecedented cluster of children presenting with MIS-C, a new hyperinflammatory syndrome in children related to the

current COVID-19 pandemic—the recognition of which led to a national alert," said the study's lead author, Shema Hameed, M.B.B.S., consultant pediatric radiologist at Evelina London Children's Hospital.

For the study, researchers performed a [retrospective review](#) of clinical, laboratory and imaging findings of the first 35 children under age 17 who were admitted to the pediatric hospital that met the case definition for MIS-C. The children were admitted from April 14 to May 9, 2020, and included 27 boys and eight girls, with a median age of 11 years old.



Thoracic findings in a 5-year-old boy with Multisystem Inflammatory Syndrome in Children (MIS-C). (a) Chest radiograph 4 days after admission shows perihilar bronchial wall cuffing and interstitial thickening with hazy surrounding airspace consolidation extending towards the peripheries. (b, c) Coronal and axial contrast-

enhanced CT chest 1 day after the radiograph shows diffuse bilateral airspace consolidation with an anteroposterior gradient suggestive of acute respiratory distress syndrome (ARDS). Credit: Radiological Society of North America

The most common clinical presentation was fever, found in 33 (94%) of the children, gastrointestinal symptoms including abdominal pain, vomiting and diarrhea in 30 (86%) of the children, rash (13 or 37%) and conjunctivitis (9 or 26%). Twenty-one children (60%) were in shock. Clinical status was severe enough to warrant management in the pediatric intensive care unit in 24 of 35 children (69%), of which 7 (20%) required mechanical ventilation and 20 (57%) inotropic support.

Two children required extracorporeal membrane oxygenation (ECMO) due to severe myocardial dysfunction. Lab tests revealed that all of the children had abnormal white blood cell counts.

The study identified a pattern of imaging findings in post COVID-19 MIS-C, including airway inflammation, rapidly progressive pulmonary edema, coronary artery aneurysms and extensive abdominal inflammatory changes within the right iliac fossa.

All 35 children underwent chest X-ray due to fever, sepsis or features of multisystem inflammation. Nineteen X-rays were abnormal, the most common finding being that of bronchial wall thickening.



15-year-old male presenting with fever, sepsis, and shock with impaired cardiac function. Coronal reconstruction of a contrast-enhanced CT chest shows multiple round consolidative foci with surrounding ground glass halos (red arrow). Credit: Radiological Society of North America

The predominant findings on chest CT were basal consolidation, or part of the lung filling with fluid; and collapsed lung with pleural effusions, or build-up of fluid in the outer membranes of the lungs.

Abdominal ultrasound findings included inflammatory changes within the right iliac fossa, with mesenteric fat stranding, lymphadenopathy and bowel wall thickening, as well as free fluid in the pelvis.

"As pediatric radiologists, we were interested in the emerging pattern of imaging findings that we observed in these children," Dr. Hameed said. "Our intention is to bring these findings to the attention of the wider radiological community."

The authors advise that future studies should include a larger group of patients, ideally utilizing multi-center databases to assess the radiological findings alongside the complex clinical course of these young patients.

More information: "Spectrum of Imaging Findings on Chest Radiograph, US, CT, and MRI Images in Multisystem Inflammatory Syndrome in Children (MIS-C) Associated with COVID-19." *Radiology* (2020)

Provided by Radiological Society of North America

Citation: New study looks at post-COVID-19 emerging disease in children (2020, June 25)
retrieved 6 May 2024 from

<https://medicalxpress.com/news/2020-06-post-covid-emerging-disease-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.