

Researchers working to improve quality of life for seniors in long-term care

June 17 2020



Credit: CC0 Public Domain

Simon Fraser University researchers are hoping their latest study on seniors will help to address one of their biggest physical challenges—injury from falls.

Falls cause more than 95 percent of hip [fractures](#) in [older adults](#) and

recovery results can be grim. About 30,000 people experience a [hip fracture](#) each year in Canada. While most will face surgery, months of recovery time and reduced independence and physical activity, others may succumb to their injuries.

"About 25 percent of older adults in [long-term care](#) who experience hip fracture die within six months, so it's really important that we direct efforts to prevent falls and also to prevent hip fracture in the event of a fall, especially in the high-risk, long-term care setting," says study lead, Steve Robinovitch, an SFU professor in the Department of Biomedical Physiology and Kinesiology.

While COVID-19 has highlighted the need to improve standards in [long-term care facilities](#), the researchers say addressing issues related to safe mobility is also key to improving seniors' quality of life.

"We need to remember that it's not just infectious disease that strikes this population, it's also problems of mobility and injuries due to falls that are often on top of their list of challenges," says Robinovitch.

Since 2007, Robinovitch has partnered with two Lower Mainland long-term care homes to record and analyze video footage of falls in common areas.

For the most recent study, published in the *Journal of Bone and Mineral Research*, the team analyzed 2,377 falls experienced by 646 residents. Only 30 of the falls caused hip fracture, a little over one percent of the total number of falls recorded.

"Only a small portion of falls caused hip fracture but the problem is that falls are so common, with the average resident in long-term care falling three times per year, that the cumulative toll is enormous," says Robinovitch.

The team found that all hip fractures were caused by falls from standing height that caused impact to the pelvis on the fracture side. While backward falls were safest, forward falls were just as likely as sideways falls to hip fracture, due to rotation during descent. Factors such as arresting the fall with outstretched hands, body weight or osteoporosis had no effect.

However, falling while wearing a hip protector reduced the risk of hip fracture by over twofold.

"Our study shows that older adults in long-term care in British Columbia are wearing and benefitting from hip protectors. Hip protectors were worn in over 70 percent of falls, and halved the risk for hip fracture. But across Canada, there's evidence that only 10 to 20 percent of people in long-term care are wearing hip protectors," Robinovitch explains.

"We're working with partners in Ontario to develop strategies to increase uptake of hip protectors in long-term care homes, and developing strategies to translate the successful B.C. model to other areas of Canada and internationally."

He says the team is continuing to analyze the data to gain a more precise understanding of the loads that are applied to bones during falls. These data can be used to help improve the design of hip protectors to provide more effective protection from the types of impacts that cause hip fractures.

The study also revealed that mobility aids reduce the likelihood of older adults fracturing a hip during a fall by over threefold. Robinovitch says families and care providers can encourage seniors to use their walkers to preserve mobility and maintain their independence while providing protection against injury.

Study co-author Fabio Feldman is the director of clinical quality and patient safety for Fraser Health. Feldman implemented a successful hip protector program in long-term care for Fraser Health, which is now being used across the province.

"This research is so important for us to better understand the characteristics of falls that result in hip fracture and shed a light on what are the possible interventions that we can have in place," he says.

Feldman, who is also an SFU alumnus, adds that he hopes their research leads to a shift from focusing on fall prevention to injury prevention.

"There are many ways that you can reduce the risk of having a fall but in the senior population, especially long-term care, quality of life is important," Feldman says. "There are different ways to make people safe without restricting their activities."

More information: Yijian Yang et al, The Effect Of Fall Biomechanics On Risk For Hip Fracture In Older Adults: A Cohort Study Of Video-Captured Falls In Long-Term Care, *Journal of Bone and Mineral Research* (2020). [DOI: 10.1002/jbmr.4048](https://doi.org/10.1002/jbmr.4048)

Provided by Simon Fraser University

Citation: Researchers working to improve quality of life for seniors in long-term care (2020, June 17) retrieved 12 September 2024 from <https://medicalxpress.com/news/2020-06-quality-life-seniors-long-term.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
