


# Silicones may lead to cell death

June 12 2020

---

RESEARCH

# Silicones may lead to cell death



Credit: Radboud University Nijmegen

Silicone molecules from breast implants can initiate processes in human cells that lead to cell death. Researchers from Radboud University have demonstrated this in a new study published on 12 June in *Scientific Reports*. "However, there are still many questions about what this could mean for the health effects of silicone breast implants. More research is therefore urgently needed," says Ger Pruijn, professor of Biomolecular Chemistry at Radboud University.

The possible side effects of silicone breast implants have been debated for decades. There are known cases where the implants have led to severe fatigue, fever, muscle and joint aches, and concentration disturbance. However, there is as yet no scientific study demonstrating the effect silicone molecules can have on human cells that could explain these side effects.

## **Silicone in the body**

It is a known fact that breast implants 'bleed', i.e. silicone molecules from the implant pass through the shell and enter the body. Earlier research, in 2016, by Dr. Rita Kappel, [plastic surgeon](#), and Radboud university [medical center](#), found that silicone molecules can then migrate through the body via the bloodstream or lymphatic system. The biochemists at Radboud University next asked themselves the follow-up question: what effect might silicone molecules have on cells exposed to it?

## Cultured cells

Experiments with cultured cells showed that silicones appeared to initiate [molecular processes](#) that lead to cell death. "We observed similarities with molecular processes related to programmed [cell death](#), a natural process called apoptosis that has an important function in clearing cells in our body. This effect appeared to depend on the dose of silicone and the size of the silicone molecules. The smaller the molecule, the stronger the effect," according to Pruijn.

To investigate the effect of silicones on human cells, the researchers have added small silicone molecules—which also occur in silicone [breast implants](#)—to three different types of cultured human cells. "One cell was more sensitive to the effect of silicones than the other two cell types. This suggests that the sensitivity of human cells to silicones varies."

## Open questions

The effects the researchers have found lead to many new questions. "We observed that silicones induce molecular changes in cells, but we don't know yet whether these changes could, for example, lead to an autoimmune response, which could in part explain the negative side effects of implants," says Pruijn.

"Caution is advised with drawing conclusions based on these findings because we used cultured cells in our research, not specific [human cells](#) such as brain cells or muscle [cells](#). Further research is required to get more clarity."

**More information:** Carla Onnekink et al, Low molecular weight silicones induce cell death in cultured cells, *Scientific Reports* (2020). [DOI: 10.1038/s41598-020-66666-7](https://doi.org/10.1038/s41598-020-66666-7)

Provided by Radboud University Nijmegen

Citation: Silicones may lead to cell death (2020, June 12) retrieved 27 April 2024 from <https://medicalxpress.com/news/2020-06-silicones-cell-death.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.