

Antidepressant does not improve post-stroke recovery

July 21 2020



Credit: CC0 Public Domain

The antidepressant fluoxetine has been suggested as a means to improve brain recovery after acute stroke. However, a large randomized study on

stroke patients at 35 Swedish hospitals shows that the drug has no such effect. The study, which was led by researchers at Karolinska Institutet, is published in *The Lancet Neurology*.

Every year almost 14 million people around the world suffer a stroke, and half of those who survive end up with a permanent functional impairment. Research on animals and small studies on humans have shown that fluoxetine, an SSRI drug that inhibits the uptake of serotonin in the brain, can promote post-stroke brain recovery. Animal studies have shown that the treatment causes new cells to be formed in the damaged area of the brain.

The effect on [functional ability](#) has now been interrogated in a large randomized study of acute [stroke patients](#) (the EFFECTS study). The researchers also studied [adverse reactions](#) to the drug and its effect on depression.

No recovery improvement

"Our study shows that fluoxetine does not improve recovery after stroke," says Erik Lundström, stroke doctor and principal investigator of the EFFECTS study and researcher at the Department of Clinical Neuroscience. "The number of depressions did indeed decrease, but the risk of bone fractures increased."

The study included 1,500 patients from 35 Swedish hospitals between October 2014 and June 2019, making it the largest ever randomized controlled stroke study in Sweden.

The patients were randomly placed in either a group that received six months' fluoxetine (20 mg) treatment or a [placebo group](#), without the participants or the researchers knowing who was assigned to which. Functional ability was then measured using the modified Rankin scale

(mRS), which is the most common scale for rating the degree of post-stroke loss of function.

Three collaborative studies

EFFECTS collaborates with two other academic-driven studies of fluoxetine treatment after stroke (FOCUS and AFFINITY). The collated results from the 6,000 or so patients included in the three studies will be presented within the year.

"My advice is to refrain from using fluoxetine as a preventative treatment following [stroke](#)," says Dr. Erik Lundström.

More information: Erik Lundström et al. Safety and efficacy of fluoxetine on functional recovery after acute stroke (EFFECTS): a randomized, double-blind, placebo-controlled trial, *The Lancet Neurology* (2020). [DOI: 10.1016/S1474-4422\(20\)30219-2](https://doi.org/10.1016/S1474-4422(20)30219-2)

Provided by Karolinska Institutet

Citation: Antidepressant does not improve post-stroke recovery (2020, July 21) retrieved 10 May 2024 from <https://medicalxpress.com/news/2020-07-antidepressant-post-stroke-recovery.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--