

Bacteria in infants' first stool may indicate their risk of obesity

July 8 2020



Credit: CC0 Public Domain

Meconium—the earliest stool of an infant—is composed of materials ingested during the time the infant spends in the uterus. A new study published in *Pediatric Obesity* found that the types of normal bacteria

found in the meconium may predict an infant's likelihood of later developing obesity.

In the study of 212 newborns, children who became overweight at 3 years of age differed in their [meconium](#) bacterial makeup from those with [normal weight](#), having a higher proportion of bacteria in the Bacteroidetes phylum (29% versus 15%).

"The concept of fetal microbiome is controversial and the colonization process after birth is better understood than the possible fetal colonization; however, there are many prenatal factors affecting the microbial composition of the baby's first stool, such as the mother's use of antibiotics during pregnancy and biodiversity of the home environment during pregnancy," said corresponding author Katja Korpela, MD, of the University of Oulu, in Finland. "It is very interesting that the microbiome formed before [birth](#) is possibly linked to a child's subsequent weight status."

More information: Katja Korpela et al, Microbiome of the first stool and overweight at age 3 years: A prospective cohort study, *Pediatric Obesity* (2020). [DOI: 10.1111/ijpo.12680](https://doi.org/10.1111/ijpo.12680)

Provided by Wiley

Citation: Bacteria in infants' first stool may indicate their risk of obesity (2020, July 8) retrieved 5 May 2024 from <https://medicalxpress.com/news/2020-07-bacteria-infants-stool-obesity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
