

Gut bacteria improve type 2 diabetes risk prediction

July 6 2020



The composition and function of bacteria in the human intestine--the so-called gut microbiome--changes as the day progresses. This was established by researchers based in Freising at ZIEL - Institute for Food & Health of the Technical University of Munich (TUM) with one of the largest studies related to microbiomes and diabetes comprising more than 4000 participants. These daily variations in the gut microbiome cease to exist in people suffering from type 2 diabetes. Credit: A. Heddergott/TUM



The microbial composition of the intestines is complex and varies widely from one individual to another. Many factors such as environmental factors, lifestyle, genetics or illnesses affect the intestinal ecosystem of helpful gut bacteria.

Dirk Haller, Professor for Nutrition and Immunology at TUM, and his team have examined the importance of daytime-dependent fluctuations of the <u>gut microbiome</u> in relation to type 2 diabetes; they present their study encompassing more than 4000 people and it is the first study in this field based on a large prospective human cohort.

The relationship between gut bacteria and medical conditions

"In order to see whether changes in the gut microbiome allow conclusions about medical conditions, so-called prospective cohort studies are required," explained Prof. Haller.

In these prospective cohort studies, a cross section of the population is being observed; however, none of the participants showed any signs of <u>disease</u>. This population is being re-examined over time. This way, researchers can find out whether a certain observation may be typical for future occurrences of diseases.

Diagnosis and outlook of type 2 diabetes may be improved

"When certain gut <u>bacteria</u> do not follow a day-night rhythm, so if their number and function does not change over the course of the day, this can be an indicator for a potential type 2 diabetes disease. Knowing this can improve diagnosis and outlook of type 2 diabetes," said Chronobiologist Dr. Silke Kiessling, another contributor to the study.



These arrhythmic bacteria—those that are not changing between day and night—are a marker for potential disease. Researchers refer to this as a risk signature. "Mathematical models also show that this microbial risk signature consisting of arrhythmic bacteria helps diagnosing diabetes," explained Sandra Reitmeier, first author on the study.

Primarily, the scientists analyzed data from an existing independent cohort by Helmholtz Zentrum München. The diabetes-related results were validated using additional cohorts from Germany. "By comparing our data to cohorts in England, we could confirm that there is—among other things—a strong regional factor affecting the microbial ecosystem. Therefore, there is a demand for finding locally specified arrhythmic risk signatures," elaborated Haller.

Nutritionist Haller emphasizes that "apart from bacteria and their variations over the course of the day, other parameters such as the body mass index play a role in being able to better predict a person's future medical conditions."

Intestinal bacteria's day and night rhythm as starting point for further research

Registering the time of day when taking human fecal samples for research purposes can heavily influence disease diagnostics. "Documenting these timestamps is essential for improving risk markers," Prof. Haller emphasizes.

This research substantiates the hypothesis that changes in the microbiome have an effect of nutrition-related diseases. How <u>gut</u> <u>bacteria</u> changing (or not changing) during the day affect other microbiome-associated diseases such as Crohn's disease or intestinal cancer may be subject to further scientific examination.



The results of this study are of particular importance for further work in the Collaborative Research Center of "Microbiome Signatures" (<u>www.sfb1371.tum.de/</u>), as <u>cohort</u> studies offer valuable possibilities of comparing data of healthy and ill subjects, particularly in the context of clinical studies.

More information: Sandra Reitmeier et al, Arrhythmic Gut Microbiome Signatures Predict Risk of Type 2 Diabetes, *Cell Host & Microbe* (2020). DOI: 10.1016/j.chom.2020.06.004

Provided by Technical University Munich

Citation: Gut bacteria improve type 2 diabetes risk prediction (2020, July 6) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2020-07-gut-bacteria-diabetes.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.