

## High levels of antibiotic-resistant bacteria found on equipment in communal gyms

July 24 2020

Research presented at ASM Microbe Online found that 43% of Staphylococcus bacteria found on exercise equipment in university gyms were ampicillin-resistant, with 73% of those isolates being resistant to multiple additional drugs. The late Xin Fan, Ph.D., and her student Chase A. Weikel of West Chester University (WCU) conducted the research in cooperation with WCU's John M. Pisciotta, Ph.D., associate professor of Biology.

According to the U.S. Centers for Disease Control and Prevention, roughly 120,000 S. aureus bacteremia cases resulted in 20,000 deaths in 2017. Skin abrasions are a common route of entry of pathogenic S. aureus strains. As highlighted by the ongoing COVID-19 pandemic, there is increasing <u>public concern</u> regarding communal areas as a bastion of infectious microorganisms.

Results of the study found 43% of 462 S. aureus isolates recovered from 45 different <u>exercise equipment</u> surfaces were ampicillin resistant. Of 60 representative ampicillin-resistant isolates, 73% were resistant to two or more additional drugs including erythromycin and sulfisoxazole.

"These results suggest regularly contacted surfaces in different recreational environments can harbor multi-drug resistant S. aureus (MDRSA) and should be disinfected frequently to best maintain <u>public</u> <u>health</u> and community wellbeing," said Chase A. Weikel, a 2018 graduate of West Chester University and current graduate student at Thomas Jefferson University in Philadelphia.



Samples were collected from 2 university recreational facilities. The surfaces gym patrons frequently touched, including dumbbells and barbell handles, cable pull grips, kettlebells, elliptical and treadmill handles were swabbed and plated on Mannitol Salt Agar (MSA). This selective and differential media was used to isolate and presumptively identify S. aureus. Isolates were replicated to MSA plus ampicillin. Isolates additionally resistant to oxacillin or penicillin were subsequently screened using CHROMagar MRSA; a sensitive and specific media used to screen for methicillin-resistant S. aureus (MRSA). Isolates that tested positive using CHROMagar were subjected to additional confirmatory methods including latex agglutination assay. Microscopy was used to confirm Gram-positive status and cellular morphology and arrangement.

This research is being presented as an ePoster at the ASM Microbe conference which is proceeding in an on-line format in August 2020. ASM Microbe Online brings you the dynamic, cutting-edge science of ASM Microbe 2020, the annual meeting of the American Society for Microbiology. Explore the latest research in the microbial sciences with ePosters, hear from experts in the field during live keynotes and access track-related content with a curated selection of on-demand sessions.

## Provided by American Society for Microbiology

Citation: High levels of antibiotic-resistant bacteria found on equipment in communal gyms (2020, July 24) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2020-07-high-antibiotic-resistant-bacteria-equipment-gyms.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.