

Higher vitamin C, carotenoid intake may lower risk for T2DM

July 9 2020



(HealthDay)—Higher intakes of vitamin C and carotenoids, as well as

total whole grains, are associated with a lower risk for type 2 diabetes, according to two studies published online July 8 in *The BMJ*.

Ju-Sheng Zheng, Ph.D., from the University of Cambridge School of Clinical Medicine in the United Kingdom, and colleagues conducted a case-cohort study involving 9,754 participants with incident type 2 [diabetes](#) and a subcohort of 13,662 individuals from the European Prospective Investigation into Cancer and Nutrition cohort. The researchers found that higher plasma vitamin C was associated with a [reduced risk](#) for developing type 2 diabetes in a multivariable adjusted model (hazard ratio per standard deviation, 0.82). A similar inverse association was seen for total carotenoids (hazard ratio per standard deviation, 0.75).

Yang Hu, Sc.D., from the Harvard T.H. Chan School of Public Health in Boston, and colleagues examined the associations between intake of total and individual whole grain foods and type 2 diabetes using data from the Nurses' Health Study, Nurses' Health Study II, and Health Professionals Follow-Up Study involving 158,259 women and 36,525 men. The researchers identified 18,629 participants with type 2 diabetes during 4,618,796 person-years of follow-up. Participants in the highest versus the lowest category of total whole grain consumption had a 29 percent lower rate of type 2 diabetes after adjustment for lifestyle and dietary risk factors.

"These findings provide further support for the current recommendations that promote increased consumption of whole grain as part of a healthy diet for the prevention of type 2 diabetes," Hu and colleagues write.

Several authors from the Hu study disclosed financial ties to the biopharmaceutical and nutrition industries.

More information: [Abstract/Full Text - Zheng](#)
[Abstract/Full Text - Hu](#)

Copyright © 2020 [HealthDay](#). All rights reserved.

Citation: Higher vitamin C, carotenoid intake may lower risk for T2DM (2020, July 9) retrieved 23 May 2024 from <https://medicalxpress.com/news/2020-07-higher-vitamin-carotenoid-intake-t2dm.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.