

## **Iodine exposure in the NICU may lead to decrease in thyroid function, study suggests**

July 7 2020

---



Credit: Pixabay/CC0 Public Domain

Exposure to iodine used for medical procedures in a neonatal intensive care unit (NICU) may increase an infant's risk for congenital hypothyroidism (loss of thyroid function), suggests a study by

researchers at the National Institutes of Health and other institutions. The authors found that infants diagnosed with congenital hypothyroidism following a NICU stay had higher blood iodine levels on average than infants who had a NICU stay but had normal thyroid function. Their study appears in the *Journal of Nutrition*.

"Limiting [iodine](#) exposure among this group of infants whenever possible may help lower the risk of losing thyroid function," said the study's first author, James L. Mills, M.D., of the Epidemiology Branch at NIH's Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD).

Congenital hypothyroidism is a partial or complete loss of thyroid function. The thyroid, located in the throat, makes iodine-containing hormones that regulate growth, brain development and the rate of chemical reactions in the body. Treatment consists of thyroid hormone therapy and must begin within four weeks after birth or permanent intellectual disability may result.

In the United States, all infants are routinely screened for the condition by collecting a small sample of blood from an infant's heel and analyzing it for thyroid stimulating hormone. Infants with a high level of thyroid stimulating hormone are referred for further testing.

To conduct the study, the researchers analyzed blood spots for their iodine content. They compared blood iodine levels from 907 children diagnosed with [congenital hypothyroidism](#) to those of 909 similar children who did not have the condition. This included 183 infants cared for in the NICU—114 of whom had congenital hypothyroidism and 69 who did not.

Overall, the researchers found no significant difference between blood iodine concentrations in those who had congenital hypothyroidism and

those in the control group. Because very high or very low iodine levels increase the risk for congenital hypothyroidism, they also looked at those infants having the highest and lowest iodine levels.

Children with congenital hypothyroidism were more likely to have been admitted to a NICU than those without congenital hypothyroidism. When the researchers considered only those infants with a NICU stay, they found that the group with congenital hypothyroidism had significantly higher iodine levels than those without the condition who also had a NICU stay. Similarly, those with congenital hypothyroidism and a NICU stay tended to have higher blood iodine than children with the condition who did not have a NICU stay.

The researchers were unable to obtain information on the [medical procedures](#) the infants may have undergone during their time in the NICU. Iodine solutions are commonly used as disinfectants to prepare the skin for surgical or other procedures. Preterm infants absorb iodine more readily through their skin than older infants. Iodine also is given internally for imaging procedures used in infants.

The researchers said that the higher iodine levels seen among infants with congenital hypothyroidism and a NICU stay may have resulted from exposure to iodine during a medical procedure. Because of this possibility, they cautioned NICU staff to use disinfectants that do not contain iodine whenever possible and to avoid exposing [infants](#) to iodine unless absolutely necessary.

Provided by NIH/Eunice Kennedy Shriver National Institute of Child Health and Human Development

Citation: Iodine exposure in the NICU may lead to decrease in thyroid function, study suggests (2020, July 7) retrieved 4 May 2024 from <https://medicalxpress.com/news/2020-07-iodine->

[exposure-nicu-decrease-thyroid.html](http://exposure-nicu-decrease-thyroid.html)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.