

Low-carbohydrate diet may be associated with lower risk of blinding eye disease

July 23 2020



Credit: Pixabay/CC0 Public Domain

Following a long-term diet that's low in carbohydrates and high in fat and protein from vegetables may lower the risk of the most common subtype of glaucoma. In a first-of-its-kind study, a researcher at New York Eye and Ear Infirmary of Mount Sinai (NYEE) helped discover that if at-risk groups adhere to these dietary restrictions, they may reduce their risk of developing primary open angle glaucoma with early

paracentral visual loss by 20 percent. Results from the research have been published in the July 22 issue of *Eye*.

The study is important because [glaucoma](#) is the leading cause of blindness in the United States and primary open angle glaucoma (POAG) is the most common type. POAG is the leading cause of optic nerve degeneration that is related to the pressure level inside the eye, but other factors also contribute to this condition. Patients typically experience few or no symptoms until the disease progresses and they have vision loss.

"A [diet](#) low in carbohydrates and higher in fats and proteins results in the generation of metabolites favorable for the mitochondrion-rich optic nerve head, which is the site of damage in POAG. This dietary pattern has already been shown to have favorable results for epilepsy and showed some promising results for Parkinson's and Alzheimer's diseases," said co-corresponding author Louis R. Pasquale, MD, FARVO, Deputy Chair for Ophthalmology Research for the Mount Sinai Health System. "It's important to note that a low-carbohydrate diet won't stop glaucoma progression if you already have it, but it may be a means to preventing glaucoma in high-risk groups. If more patients in these high-risk categories—including those with a family history of glaucoma—adhered to this diet, there might be fewer cases of vision loss."

Past studies have shown that a [ketogenic diet](#) (low-carbohydrate and high-fat) has a protective effect against neurologic disorders. Ketone bodies (energy compounds produced as the body metabolizes fats) are substituted for glucose as a major energy source for the brain and using more of these under a ketogenic diet may improve function and slow down neuronal degeneration. However, more recent studies have shown that a low-carbohydrate diet that does not restrict protein or total calories may have similar neuroprotective properties; this diet has been

recognized as a more practical alternative to a ketogenic diet, as it's easier to follow and does not have the adverse effects of the ketogenic diet (which range from headache, weakness, and irritability to constipation, nausea, and vomiting).

A team of researchers wanted to know if a low-carbohydrate diet could positively impact the optic nerve. The optic nerve sits in the back of the eye and transfers visual information from the retina to the brain through electrical impulses. The optic nerve has a large concentration of mitochondria (mitochondria represent the major source of a cell's energy supply and lead to a cell's survival), and has high-energy requirements. Since glaucoma is a condition that may be associated with [mitochondrial dysfunction](#), researchers wanted to find out if substituting protein and fat for carbohydrates in the diet would enhance mitochondrial activity, maintain optic nerve function, and prevent optic nerve degeneration in this blinding eye disease. They performed a large-scale meta-analysis to get this answer.

They followed 185,000 adult participants from three large studies in the United States, conducted between 1976 and 2017. Participants were female nurses and male health professionals between the ages of 40 and 75. Every two to four years, they filled out food frequency questionnaires that assessed what they ate and drank. They also answered questions about their health and what diseases, if any, they might be developing. If they said they had glaucoma, the researchers asked their treating eye care providers to send medical records to determine if they had POAG.

The research team created statistical models based on the patients' questionnaire responses, dividing them into groups based on carbohydrate intake, so they could look across the spectrum from high to low carbohydrate intake and see any possible relationship with POAG. They specifically looked at three different ways of achieving a low-

carbohydrate diet: substituting animal-based fats and proteins for carbohydrates; substituting plant-based based fats and proteins for carbohydrates; and replacing carbohydrates with high fats and proteins regardless of the source. Researchers then calculated the relative risk of POAG after adjusting for multiple factors for each of the dietary patterns including age, race, and body mass index.

Patients in the low-carbohydrate intake group who followed a diet of increased plant-based fat and protein were associated with a 20 percent lower risk of developing POAG subtype with paracentral visual field loss compared to those in the high-carbohydrate intake group. However, the researchers did not find any association between POAG and a low-carbohydrate diet without accounting for the source protein or fat, and they did not find any association between glaucoma and an animal-based low-carbohydrate diet. Their findings suggest vegetable sources may be more beneficial than animal sources for a low-carbohydrate diet with respect to reducing risk of the specific glaucoma subtype with early paracentral visual loss.

"This was an observational study and not a clinical trial, so more work is needed as this is the first study looking at this dietary pattern in relation to POAG. The next step is to use artificial intelligence to objectively quantify paracentral visual loss in our glaucoma cases and repeat the analysis," adds Dr. Pasquale. "It's also important to identify patients who have a genetic makeup of primary open angle glaucoma who may benefit from a [low-carbohydrate diet](#). This dietary pattern may be protective only in people with a certain genetic makeup."

More information: Akiko Hanyuda et al. Low-carbohydrate-diet scores and the risk of primary open-angle glaucoma: data from three US cohorts, *Eye* (2020). [DOI: 10.1038/s41433-020-0820-5](https://doi.org/10.1038/s41433-020-0820-5)

Provided by The Mount Sinai Hospital

Citation: Low-carbohydrate diet may be associated with lower risk of blinding eye disease (2020, July 23) retrieved 25 April 2024 from <https://medicalxpress.com/news/2020-07-low-carbohydrate-diet-eye-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.