

Middle aged face more years of ill health than baby boomers

July 15 2020, by Mark Greaves



Credit: [Mabel Amber](#) from [Pixabay](#)

Despite expecting to live longer, people in their 40s and 50s are likely to suffer more years of ill health than older generations now in their 60s and early 70s, according to a new UCL-led study.

The study, published in the journal *Population Studies*, compared generations born between 1945 and 1980 and found a greater prevalence of ill health among those born later, with these younger cohorts more likely to rate their health as poor and have clinically measured poor health at equivalent ages during their working lives.

The researchers concluded that, although life expectancy has increased in recent decades, many of the years gained are likely to be spent in poor health, with conditions such as [diabetes](#) and obesity affecting people earlier.

Lead author, Dr. Stephen Jivraj (UCL Epidemiology & Public Health), said: "Our study shows that, for those born between 1945 and 1980, the overall trend is towards an increasing proportion of years in poor health, with some health conditions beginning at an earlier age. This has worrying implications for healthcare services, which already face increased demand because of an aging population."

Researchers looked at data from 135,189 people aged between 25 and 64 who took part in the Health Survey for England (HSE), an annual household survey with a nationally representative sample, between the years of 1991 and 2014.

Participants were asked whether they had poor health, a long-term illness, and a range of chronic conditions such as cardiovascular disease and diabetes. Nurses also took objective measurements of hypertension, body mass index (BMI) and glycated hemoglobin (clinical diabetes).

Researchers compared the results for different age groups and used the data to calculate changes in healthy life expectancy over the generations as well as years likely to be spent in poor health.

They calculated that half of the gains in life expectancy between 1993

and 2003 would likely be spent in poor health, falling to a fifth of the gains between 2003 and 2013. Later-born cohorts were more likely to have diabetes, to be overweight and to report having cardiovascular disease and [poor health](#) in general. Later-born men were more likely to report high blood pressure.

Senior author, Professor George Ploubidis (UCL Center for Longitudinal Studies), said: "Earlier in the 20th century, a rise in [life expectancy](#) went hand in hand with an increase in healthy lifespan—younger generations were living longer, healthier lives.

"It appears that, for those generations born between 1945 and 1980, this trend has stalled. Those born later are expected to live longer on average, but with more years of ill [health](#)."

More information: Stephen Jivraj et al. Living longer but not necessarily healthier: The joint progress of health and mortality in the working-age population of England, *Population Studies* (2020). [DOI: 10.1080/00324728.2020.1767297](#)

Provided by University College London

Citation: Middle aged face more years of ill health than baby boomers (2020, July 15) retrieved 27 April 2024 from <https://medicalxpress.com/news/2020-07-middle-aged-years-ill-health.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.