

Office-based exercise could reduce heart disease risk by 15%

July 14 2020



Dr Niels Vollaard. Credit: University of Stirling

Employers could help staff reduce their risk of heart disease by 15 percent if they introduced specialised exercise bikes to the workplace, according to new research.

The University of Stirling-led study found that staff who cycle for just 18 minutes per week could realise [health](#) benefits—which also include improved fitness and reduced risk of type 2 diabetes.

As part of the research, exercise bikes were introduced to office settings and volunteers were asked to cycle for 8 minutes 40 seconds twice a week—a fraction of the 150 minutes of moderate exercise recommended under government guidelines. The novel exercise routine—known as reduced-exertion high-intensity interval training (REHIT) – allows participants to wear their work clothes and avoids the need for a post-workout shower, due to the low sweat response.

Dr. Niels Vollaard, of the University's Faculty of Health Sciences and Sport, led the study—conducted prior to the COVID-19 pandemic—alongside colleagues from Swansea University and the University of Cairo.

Dr. Vollaard said: "Many people do not perform enough exercise and, therefore, are at increased risk of developing diseases including [heart disease](#) and type 2 diabetes. One of the most commonly reported reasons for not meeting physical activity recommendations is a lack of time—with people leading busy lives, spending long working days sitting at desks, and commuting by car.

"Our team have spent the last eight years looking at alternative exercise routines to help people reap the [health benefits](#) of exercise in less time. REHIT involves easy pedalling on a stationary bike, interspersed with two short bursts of high-intensity cycling. It is time efficient and perceived as being manageable by our research participants.

"Prior to this study, this approach had only been tested in laboratories—however, we have now confirmed that it can be successfully implemented in an office setting."

Twenty-five previously inactive office employees from local authority offices in Stirling and Swansea were recruited for the research. Thirteen of the volunteers were allocated to the exercise group, while the others formed the control group, who continued with their regular lifestyle.

The research used CAR.O.L bikes, which differ to regular gym bikes by enabling users to reach their own maximum exercise intensity for very short durations. They were set up in the workplace, out of view from colleagues, and participants completed two sessions a week for six weeks. Each session lasted for 8 minutes and 40 seconds and combined easy pedalling with two short bursts of high-intensity cycling.

Maximal oxygen uptake (VO₂max) – a measurement of the maximum amount of oxygen a person can utilise during intense exercise—was recorded at the beginning and at the end of the six weeks. Importantly, VO₂max is the best measures of a person's general health and their future risk of developing heart disease or type 2 diabetes. The team also collected information on the volunteers' thoughts and feelings towards the exercise routine.

Health benefits

"We found that the REHIT routine was effective at improving the [general health](#) of the research participants," Dr. Vollaard explained. "VO₂max increased by around 10 percent, compared to the control group, which equates to a risk reduction for getting heart disease in later life of 15 percent.

"Based on previous research, we would expect that continuing with the

REHIT routine would improve VO₂max further—again, further reducing the risk of heart disease.

"We also found that participants considered the routine achievable, acceptable and enjoyable."

Due to the COVID-19 pandemic, the research team acknowledge that employers may be reluctant to introduce workplace exercise in the immediate future—and any move to do so would have to meet hygiene and social distancing guidance. However, they believe their findings provide an important insight into the possibilities that could be considered as workplaces return to 'normal'.

The paper, "Time-efficient and computer-guided sprint interval exercise training for improving health in the workplace: a randomised mixed-methods feasibility study in office-based employees," is published in *BMC Public Health*.

More information: Richard S. Metcalfe et al. Time-efficient and computer-guided sprint interval exercise training for improving health in the workplace: a randomised mixed-methods feasibility study in office-based employees, *BMC Public Health* (2020). [DOI: 10.1186/s12889-020-8444-z](https://doi.org/10.1186/s12889-020-8444-z)

Provided by University of Stirling

Citation: Office-based exercise could reduce heart disease risk by 15% (2020, July 14) retrieved 25 April 2024 from <https://medicalxpress.com/news/2020-07-office-based-heart-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.