

Pandemic caused 18 pc rise in deaths in US: study

July 1 2020



This scanning electron microscope image shows SARS-CoV-2 (yellow)—also known as 2019-nCoV, the virus that causes COVID-19—isolated from a patient, emerging from the surface of cells (blue/pink) cultured in the lab. Credit: NIAID-RML

The coronavirus pandemic in the US claimed at least 122,000 more lives

than would be expected in a normal year, for a rise of 18 percent, says a study released Wednesday.

But this is just a national average, and the excess [death](#) rate was particularly high in virus hot spots such as New York City, which buried three times more people than usual and up to seven times as many during the peak of the pandemic, according to a week by week study carried out at Yale University and published in the journal *JAMA Internal Medicine*.

In New York City, the expected deaths under a demographic model based on statistics from previous years would be 13,000 from March 1 through the end of May. But this time the number of deaths recorded was 38,170.

What is more, throughout the first phase of the pandemic in the US the official COVID-19 death toll was widely underestimated, the statistics in this study show.

The total number of extra deaths was far greater than that of fatalities officially blamed on the coronavirus. This is because many people who died were not tested for the virus, or because the way death certificates are filled out is not standardized in the US. So 22 percent of the above-normal deaths had no official link to the coronavirus.

States such as Texas and Arizona, which went relatively unscathed in the spring—but are now hit hard in a new virus surge—were the worst off by this measure. More than half of the excess deaths went unexplained, with no official link to the pandemic.

But this margin got smaller as more testing was carried out in the US.

"The gap between the official COVID-19 tally and the excess deaths has been shrinking over time and has nearly disappeared in some places, like

New York City," Daniel Weinberger, an associate professor of epidemiology at the Yale School of Public Health and first author of the study, told AFP.

"How reliably the official tolls capture the full burden of excess deaths still varies considerably between states," he added.

The official COVID-19 death toll is relatively reliable in New York, Massachusetts or Minnesota, for instance, the study shows.

The study does not address the issue of deaths caused indirectly by the [pandemic](#). These are people who died of other causes, such as a [heart attack](#) or stroke and refused to go to a hospital for fear of getting infected with the [coronavirus](#).

Separate data show that these causes of death increased although Weinberger said he does not think they contributed a lot to the overall excess deaths.

More information: Daniel M. Weinberger et al. Estimation of Excess Deaths Associated With the COVID-19 Pandemic in the United States, March to May 2020, *JAMA Internal Medicine* (2020). [DOI: 10.1001/jamainternmed.2020.3391](#)

© 2020 AFP

Citation: Pandemic caused 18 pc rise in deaths in US: study (2020, July 1) retrieved 3 May 2024 from <https://medicalxpress.com/news/2020-07-pandemic-pc-deaths.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
