

Pimavanserin reduced symptoms of dementia-related psychosis in phase 3 trial

July 29 2020

A large-scale trial has found a treatment with pimavanserin substantially reduced psychotic symptoms and reduced risk of relapse of those symptoms compared to placebo in people with dementia, a condition for which no treatments are currently licensed.

New data presented at the Alzheimer's Association International Conference indicates that pimavanserin leads to a robust reduction in the severity of psychosis symptoms during the 12 week open-label phase of the study, regardless of the underlying dementia subtype or the severity of participants' dementia.

The international phase three relapse-prevention trial was conducted in collaboration with the University of Exeter, working with Acadia Pharmaceuticals. The trial enrolled 392 participants who were experiencing hallucinations and delusions as a result of their dementia.

Up to half of the 45 million people worldwide who are living with Alzheimer's disease will experience psychotic episodes, a figure that is even higher in some other forms of dementia. Psychosis is linked to a faster deterioration in dementia, and is disturbing for the individual and carers.

Currently there is no approved safe and [effective treatment](#) for these distressing symptoms associated with dementia-related psychosis. In people with dementia, widely-used antipsychotics lead to sedation, falls, and can double the speed at which [brain function](#) deteriorates.

Pimavanserin works differently, by blocking a very specific nerve receptor (5HT2A) in the brain. Now, it has been found to effectively reduce symptoms of psychosis in people with Alzheimer's disease and other dementias, without evidence of some of the common side effects observed with other antipsychotics.

Clive Ballard, Professor of Age-Related Diseases at the University of Exeter Medical School, who co-authored the research presented at AAIC, said: "Psychosis is a common and extremely disturbing [symptom](#) which has a severe impact on the lives of people affected by [dementia](#). We urgently need treatments that work, and I'm excited by the potential of pimavanserin to make a real difference."

Provided by University of Exeter

Citation: Pimavanserin reduced symptoms of dementia-related psychosis in phase 3 trial (2020, July 29) retrieved 5 May 2024 from <https://medicalxpress.com/news/2020-07-pimavanserin-symptoms-dementia-related-psychosis-phase.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|