


Prostate cancer: Analytics suggest when physicians should treat

July 20 2020



Credit: Pixabay/CC0 Public Domain

Using the first results from an analysis of the world's largest active surveillance prostate cancer database, the GAP3 consortium has begun to identify which patients are at risk of developing the disease and which patients can safely delay treatment. This work is presented at the virtual European Association of Urology congress.

Lead researcher Dr. Mieke Van Hemelrijck (King's College London) says, "Current methods of deciding whether or not to recommend treatment are not reliable. Our analysis shows that we should be able to produce a single global methodology that will give accurate estimates on how aggressive these cancers are. These will feed directly into the treatment decision, and give men the reassurance they need to decide on treatment."

Prostate cancer is one of the leading causes of death in men, but many men who discover they have [prostate cancer](#) have [low-risk prostate cancer](#). Over the past 10 years, an increasing number of these men have been given the option of going on [active surveillance](#) rather than being immediately treated. Active [surveillance](#) means that men continue to be monitored and tested (via PSA levels, biopsy, and other tests), with treatment only starting when the cancer shows signs of developing. The number of men on active surveillance varies from country to country, with up to 80% of men delaying treatment in some countries. However, there are no generally accepted ways of understanding who is at risk, and as many as 38% of men who start active surveillance drop out within five years.

Van Hemelrijck said, "Prostate cancer treatment can have significant side-effects such as erectile dysfunction and incontinence, so often, avoiding intrusive surgery or radiotherapy can benefit the patient. Nevertheless, being told you have cancer puts great psychological pressure on men to agree to treatment, so understanding just how aggressive the cancer is before deciding on treatment is essential. At the moment we just don't have that reassurance."

Although active surveillance is considered a real step forward in management of low risk prostate cancer, there is surprisingly little agreement on which men will benefit. Doctors consider a range of factors such as age, PSA score, biopsy details, technical details of the

[cancer](#), and so on. But the decision on whether or not to start treatment is still often subjective. The Erasmus MC Department of Urology was tasked by Movember to coordinate the development of a global database on active surveillance (the GAP3 Consortium). Dr. Van Hemelrijck worked with a team of researchers from the GAP3 Consortium to develop the world's most accurate active surveillance nomogram.

A nomogram is a treatment calculator: You feed in the details and it gives you advice on whether or not to treat. Local nomograms exist, but a global version is needed to be generally applicable. Working with data from the 14,380 patients on the Movember database (the world's largest), they were able to input data such as age, size and condition of the tumor, PSA, biopsy details, time on active surveillance, genetic factors, etc.

"Not surprisingly, we have found that even accounting for these factors there was still differences in outcomes between participating centers. But this work has shown that it will be possible to produce a nomogram which can guide treatment. Just as importantly, the work shows which additional factors need to be included in the nomogram in future to enable us to eliminate this variation and produce accurate estimates of tumor aggressiveness."

EAU Adjunct Secretary General Professor Hendrik Van Poppel (University of Leuven, Belgium), who was not involved in the study, said, "This work shows that it should be possible to develop a global nomogram—in other words, a system which allows us to predict whether active surveillance will be suitable for individual low and intermediate risk [prostate cancer](#) patients. This would be an important step forward in terms of the reassurance we can offer patients, and in choosing treatment pathways. The urology community would welcome this, and will be happy to cooperate in taking this project forward."

Provided by European Association of Urology

Citation: Prostate cancer: Analytics suggest when physicians should treat (2020, July 20)
retrieved 3 May 2024 from

<https://medicalxpress.com/news/2020-07-prostate-cancer-analytics-physicians.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.